

WEST Search History

DATE: Tuesday, June 25, 2002

<u>Set Name</u> side by side	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u> result set
<i>DB=USPT,PGPB,JPAB,EPAB,DWPI; PLUR=YES; OP=OR</i>			
L4	YKL adj 39	2	L4
L3	YKL39	1	L3
L2	L1 and YKL\$4	1	L2
L1	(verheijden)[IN] OR (boots)[IN]	380	L1

END OF SEARCH HISTORY

(FILE 'HOME' ENTERED AT 09:46:22 ON 25 JUN 2002)

FILE 'MEDLINE, CAPLUS, EMBASE, BIOSIS' ENTERED AT 09:46:36 ON 25 JUN 2002

L1 0 S PTLASAETT OR HSPTLASAETTVG
L2 211 S YKL
L3 69 S L2 AND CHONDROCYTE?
L4 42 S L3 AND (AUTOIMMUN? OR RA OR ARTHRITIS)
L5 21 DUP REM L4 (21 DUPLICATES REMOVED)
L6 19 S YKL (1N) 39
L7 10 DUP REM L6 (9 DUPLICATES REMOVED)

WU 2001 029081
WU 2000 005254
WU 2000 004917
WU 97 400684
97 00770

imm 199 96(4) 586-
J. Biol Chem 199 271 (32) 1925-20
5736507

J Biol Chem 196 271 (32) 1925-20

ile medline caplus embase biosis
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
0.21	0.21

FULL ESTIMATED COST

FILE 'MEDLINE' ENTERED AT 22:43:34 ON 24 JUN 2002

FILE 'CAPLUS' ENTERED AT 22:43:34 ON 24 JUN 2002
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'EMBASE' ENTERED AT 22:43:34 ON 24 JUN 2002
COPYRIGHT (C) 2002 Elsevier Science B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 22:43:34 ON 24 JUN 2002
COPYRIGHT (C) 2002 BIOLOGICAL ABSTRACTS INC.(R)

=> s verheijden G?/au or Boots A?/au
L1 121 VERHEIJDEN G?/AU OR BOOTS A?/AU

=> s l1 and peptide?
L2 39 L1 AND PEPTIDE?

=> s l2 and autoimmun?
L3 14 L2 AND AUTOIMMUN?

=> dup rem l3
PROCESSING COMPLETED FOR L3
L4 10 DUP REM L3 (4 DUPLICATES REMOVED)

=> dis l4 1-10 ibib abs kwic

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:sssptal644axd

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 Jan 25 BLAST(R) searching in REGISTRY available in STN on the Web
NEWS 3 Jan 29 FSTA has been reloaded and moves to weekly updates
NEWS 4 Feb 01 DKILIT now produced by FIZ Karlsruhe and has a new update
frequency
NEWS 5 Feb 19 Access via Tymnet and SprintNet Eliminated Effective 3/31/02
NEWS 6 Mar 08 Gene Names now available in BIOSIS
NEWS 7 Mar 22 TOXLIT no longer available
NEWS 8 Mar 22 TRCTHERMO no longer available
NEWS 9 Mar 28 US Provisional Priorities searched with P in CA/Caplus
and USPATFULL
NEWS 10 Mar 28 LIPINSKI/CALC added for property searching in REGISTRY
NEWS 11 Apr 02 PAPERCHEM no longer available on STN. Use PAPERCHEM2 instead.
NEWS 12 Apr 08 "Ask CAS" for self-help around the clock
NEWS 13 Apr 09 BEILSTEIN: Reload and Implementation of a New Subject Area
NEWS 14 Apr 09 ZDB will be removed from STN
NEWS 15 Apr 19 US Patent Applications available in IFICDB, IFIPAT, and IFIUDB
NEWS 16 Apr 22 Records from IP.com available in CAPLUS, HCAPLUS, and ZCAPLUS
NEWS 17 Apr 22 BIOSIS Gene Names now available in TOXCENTER
NEWS 18 Apr 22 Federal Research in Progress (FEDRIP) now available
NEWS 19 Jun 03 New e-mail delivery for search results now available
NEWS 20 Jun 10 MEDLINE Reload
NEWS 21 Jun 10 PCTFULL has been reloaded

NEWS EXPRESS February 1 CURRENT WINDOWS VERSION IS V6.0d,
CURRENT MACINTOSH VERSION IS V6.0a(ENG) AND V6.0Ja(JP),
AND CURRENT DISCOVER FILE IS DATED 05 FEBRUARY 2002
NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that
specific topic.

All use of STN is subject to the provisions of the STN Customer
agreement. Please note that this agreement limits use to scientific
research. Use for software development or design or implementation
of commercial gateways or other similar uses is prohibited and may
result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 22:43:19 ON 24 JUN 2002

=> file medline caplus embase biosis		
COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'MEDLINE' ENTERED AT 22:43:34 ON 24 JUN 2002

FILE 'CAPLUS' ENTERED AT 22:43:34 ON 24 JUN 2002
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'EMBASE' ENTERED AT 22:43:34 ON 24 JUN 2002
COPYRIGHT (C) 2002 Elsevier Science B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 22:43:34 ON 24 JUN 2002
COPYRIGHT (C) 2002 BIOLOGICAL ABSTRACTS INC.(R)

=> s verheijden G?/au or Boots A?/au
L1 121 VERHEIJDEN G?/AU OR BOOTS A?/AU

=> s l1 and peptide?
L2 39 L1 AND PEPTIDE?

=> s l2 and autoimmune?
L3 14 L2 AND AUTOIMMUN?

=> dup rem l3
PROCESSING COMPLETED FOR L3
L4 10 DUP REM L3 (4 DUPLICATES REMOVED)

=> dis l4 1-10 ibib abs kwic

L4 ANSWER 1 OF 10 CAPLUS COPYRIGHT 2002 ACS
ACCESSION NUMBER: 2001:300756 CAPLUS
DOCUMENT NUMBER: 134:320857
TITLE: Modified **peptides** and peptidomimetics for
use in immunotherapy
INVENTOR(S): Van Staveren, Catherina Joanna; Timmers, Cornelis
Marius; Van Galen, Philippus Johannes Marie; Knegtel,
Rnaldus Marcellus Alphonsus; **Boots, Anna Maria**
Helena; Miltenburg, Andreas Martinus Maria
PATENT ASSIGNEE(S): Akzo Nobel N.V., Neth.
SOURCE: PCT Int. Appl., 52 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001029081	A1	20010426	WO 2000-EP10230	20001012
W: AE, AG, AL, AU, BA, BB, BG, BR, BZ, CA, CN, CR, CU, CZ, DM, DZ, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MA, MG, MK, MN, MX, MZ, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RN: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
PRIORITY APPLN. INFO.:			EP 1999-203427 A 19991018	
OTHER SOURCE(S): MARPAT 134.320857				
AB	The invention relates to a modified peptide derived from formula I peptide H-Arg-Ser-Phe-Thr-Leu-Ala-Ser-Ser-Glu-Thr-Gly-Val-Gly-OH (peptide (263-275) of cartilage-derived protein human cartilage gp-39 (HC gp-39)) having general formula (II): Q-A1-A2-A3-A4-A5-A6-A7-A8-A9-A10-A11-A12-A13-Z. In general formula (II), A1 through A13 correspond with the amino acids of formula (I), Q corresponds with H and Z corresponds with OH. The modifications according to the present invention are selected from one or more of the groups a, b or c, consisting of (a) substitution of 1-6, preferably 1-4 amino acids at A1 through A13 with non-natural amino acids or .beta. amino acids; (b) substitution of one or more amide bonds with reduced amide bonds or ethylene isosteres; (c) substitutions at Q and/or Z and, optionally, (d) substitution of natural amino acids up to a total of 6 modifications. The peptides can be used for inducing tolerance induction in patients suffering from autoimmune diseases. The most potent compds. were Ac-Arg-Ser-Phe-Thr-Leu-Ala-Ser-Ser-Glu-Thr-Gly-Val-Gly-OH, Ac-Arg-Ser-Phe-Thr-Leu-Ala-Ser-Ser-Glu-Thr-Gly-Val-.psi.[CH2NH]-Gly-NH2, Ac-Arg-NhSer-Phe-Thr-Leu-Ala-Ser-Ser-Glu-Thr-Gly-Val-Gly-NH2 and Ac-Arg-NhSer-Phe-Thr-Leu-Ala-Ser-Ser-Glu-Thr-Gly-Val-.psi.[CH2NH]-Gly-NH2.			
REFERENCE COUNT:	5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT			
TI	Modified peptides and peptidomimetics for use in immunotherapy			
IN	Van Staveren, Catherina Joanna; Timmers, Cornelis Marius; Van Galen, Philippus Johannes Marie; Knegtel, Rnaldus Marcellus Alphonsus; Boots, Anna Maria Helena; Miltenburg, Andreas Martinus Maria			
AB	The invention relates to a modified peptide derived from formula I peptide H-Arg-Ser-Phe-Thr-Leu-Ala-Ser-Ser-Glu-Thr-Gly-Val-Gly-OH (peptide (263-275) of cartilage-derived protein human cartilage gp-39 (HC gp-39)) having general formula (II): Q-A1-A2-A3-A4-A5-A6-A7-A8-A9-A10-A11-A12-A13-Z. In general formula (II), A1 through A13 correspond with the amino acids of formula (I), Q corresponds with H and Z corresponds with OH. The modifications according to the present invention are selected from one or more of the groups a, b or c, consisting of (a) substitution of 1-6, preferably 1-4 amino acids at A1 through A13 with non-natural amino acids or .beta. amino acids; (b) substitution of one or more amide bonds with reduced amide bonds or ethylene isosteres; (c) substitutions at Q and/or Z and, optionally, (d) substitution of natural amino acids up to a total of 6 modifications. The peptides can be used for inducing tolerance induction in patients suffering from autoimmune diseases. The most potent compds. were Ac-Arg-Ser-Phe-Thr-Leu-Ala-Ser-Ser-Glu-Thr-Gly-Val-Gly-OH, Ac-Arg-Ser-Phe-Thr-Leu-Ala-Ser-Ser-Glu-Thr-Gly-Val-.psi.[CH2NH]-Gly-NH2, Ac-Arg-NhSer-Phe-Thr-Leu-Ala-Ser-Ser-Glu-Thr-Gly-Val-Gly-NH2 and Ac-Arg-NhSer-Phe-Thr-Leu-Ala-Ser-Ser-Glu-Thr-Gly-Val-.psi.[CH2NH]-Gly-NH2.			
ST	peptide peptidomimetic immunotherapy; human cartilage glycoprotein 39 peptide immune tolerance; autoimmune disease immune tolerance induction peptide			
IT	Antigens			
IT	RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)			
IT	(autoantigens, induction of specific T-cell tolerance to; modified peptides and peptidomimetics based on peptide from human cartilage glycoprotein 39 for use in immunotherapy)			
IT	Analysis			
IT	(biochem., diagnostic compn. contg. peptide and detection agent for; modified peptides and peptidomimetics based on peptide from human cartilage glycoprotein 39 for use in immunotherapy)			
IT	Allergy			
IT	(delayed hypersensitivity; modified peptides and peptidomimetics based on peptide from human cartilage glycoprotein 39 for use in immunotherapy)			
IT	Rheumatoid arthritis			
IT	(induction of specific T-cell tolerance to autoantigen of; modified peptides and peptidomimetics based on peptide from human cartilage glycoprotein 39 for use in immunotherapy)			
IT	Autoimmune disease			
IT	Diagnosis			
IT	Drug delivery systems			
IT	Immune tolerance			
IT	Immunotherapy			
IT	Peptidomimetics			
IT	(modified peptides and peptidomimetics based on peptide from human cartilage glycoprotein 39 for use in immunotherapy)			
IT	Peptides , biological studies			
IT	RL: ARG (Analytical reagent use); BPR (Biological process); BSU (Biological study, unclassified); THU (Therapeutic use); ANST (Analytical study); BIOL (Biological study); PROC (Process); USES (Uses)			
IT	(modified peptides and peptidomimetics based on peptide from human cartilage glycoprotein 39 for use in immunotherapy)			
IT	T cell (lymphocyte)			
IT	(tolerance to autoantigen; modified peptides and peptidomimetics based on peptide from human cartilage glycoprotein 39 for use in immunotherapy)			
IT	335598-47-5P	335598-48-6P	335598-49-7P	335598-50-0P
IT	335598-52-2P	335598-54-4P	335598-55-5P	335598-56-6P
IT	335598-58-8P	335598-60-2P		
IT	RL: ARG (Analytical reagent use); BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); THU (Therapeutic use); ANST (Analytical study); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)			
IT	(modified peptides and peptidomimetics based on peptide from human cartilage glycoprotein 39 for use in immunotherapy)			
IT	335598-53-3	335598-59-9		
IT	RL: ARG (Analytical reagent use); BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study,			

unclassified); PRP (Properties); THU (Therapeutic use); ANST (Analytical study); BIOL (Biological study); PROC (Process); USES (Uses)
(modified **peptides** and peptidomimetics based on
peptide from human cartilage glycoprotein 39 for use in immunotherapy)

IT 335598-46-4P
RL: ARG (Analytical reagent use); BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); ANST (Analytical study); BIOL (Biological study); PREP (Preparation); PROC (Process); RACT (Reactant or reagent); USES (Uses)
(modified **peptides** and peptidomimetics based on
peptide from human cartilage glycoprotein 39 for use in immunotherapy)

IT 335598-72-6 335598-73-7 335598-74-8
RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
(modified **peptides** and peptidomimetics based on
peptide from human cartilage glycoprotein 39 for use in immunotherapy)

IT 335598-69-1 335598-70-4 335598-71-5
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)
(modified **peptides** and peptidomimetics based on
peptide from human cartilage glycoprotein 39 for use in immunotherapy)

IT 335598-61-3D, conjugates with PAC-PEG-PS resin
RL: PRP (Properties); RCT (Reactant); RACT (Reactant or reagent)
(modified **peptides** and peptidomimetics based on
peptide from human cartilage glycoprotein 39 for use in immunotherapy)

IT 335598-62-4DP, conjugates with PAL-PEG-PS resin 335598-63-5P
335598-64-6P 335598-65-7P 335598-66-8DP, conjugates with PAL-PEG-PS resin 335598-68-0DP, conjugates with PAL-PEG-PS resin
RL: PRP (Properties); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(modified **peptides** and peptidomimetics based on
peptide from human cartilage glycoprotein 39 for use in immunotherapy)

IT 75-65-0, reactions 298-12-4, Glyoxylic acid 2462-31-9 24424-99-5
77987-49-6 88574-06-5 97807-17-5 156939-64-9
RL: RCT (Reactant); RACT (Reactant or reagent)
(modified **peptides** and peptidomimetics based on
peptide from human cartilage glycoprotein 39 for use in immunotherapy)

IT 141743-30-8P 258332-54-6P 335598-67-9P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(modified **peptides** and peptidomimetics based on
peptide from human cartilage glycoprotein 39 for use in immunotherapy)

IT 199275-21-3P
RL: SPN (Synthetic preparation); PREP (Preparation)
(modified **peptides** and peptidomimetics based on
peptide from human cartilage glycoprotein 39 for use in immunotherapy)

L4 ANSWER 2 OF 10 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 2000:84838 CAPLUS
DOCUMENT NUMBER: 132:121464
TITLE: Novel **peptides** for use in immunotherapy of autoimmune diseases
INVENTOR(S): Verheijden, Gijsbertus Franciscus Maria; Boots, Anna Maria Helena
PATENT ASSIGNER(S): Akzo Nobel N.V., Neth.
SOURCE: PCT Int. Appl., 26 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000005254	A2	20000203	WO 1999-EP5050	19990716
WO 2000005254	A3	20000615		
W: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
AU 9954112	A1	20000214	AU 1999-54112	19990716
BR 9912378	A	20010417	BR 1999-12378	19990716
EP 1100823	A2	20010523	EP 1999-940012	19990716
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
NO 2001000355	A	20010320	NO 2001-355	20010122
PRIORITY APPLN. INFO.: EP 1998-202470 A 19980723				
WO 1999-EP5050 W 19990716				

AB The invention relates to the use of novel **peptides** in a **peptide** induced tolerance therapy to prevent autoimmune disorders and in particular their use in treatment of chronic destruction of articular cartilage. The invention furthermore embraces pharmaceutical compns. comprising said **peptides** and a diagnostic method for the detection of autoreactive T cells in a test sample.

TI Novel **peptides** for use in immunotherapy of autoimmune diseases

IN Verheijden, Gijsbertus Franciscus Maria; Boots, Anna Maria Helena

AB The invention relates to the use of novel **peptides** in a **peptide** induced tolerance therapy to prevent autoimmune disorders and in particular their use in treatment of chronic destruction of articular cartilage. The invention furthermore embraces pharmaceutical compns. comprising said **peptides** and a diagnostic method for the detection of autoreactive T cells in a test sample.

ST autoimmune articular cartilage destruction tolerance immunotherapy; autoantigen tolerance autoimmune articular cartilage destruction

IT Histocompatibility antigens
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL

(Biological study); PROC (Process)
 (HLA-DR, DRB1*0401; autoantigenic **peptides** for inducing tolerance for immunotherapy of **autoimmune** diseases)

IT Cartilage
 (articular, chronic destruction; autoantigenic **peptides** for inducing tolerance for immunotherapy of **autoimmune** diseases)

IT **Autoimmune** disease
 Immunotherapy
 Protein sequences
 Rheumatoid arthritis
 (autoantigenic **peptides** for inducing tolerance for immunotherapy of **autoimmune** diseases)

IT Antigens
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (autoantigens; autoantigenic **peptides** for inducing tolerance for immunotherapy of **autoimmune** diseases)

IT Drug delivery systems
 (carriers; autoantigenic **peptides** for inducing tolerance for immunotherapy of **autoimmune** diseases)

IT Musculoskeletal diseases
 (cartilage, articular; autoantigenic **peptides** for inducing tolerance for immunotherapy of **autoimmune** diseases)

IT Cartilage
 (disease, articular; autoantigenic **peptides** for inducing tolerance for immunotherapy of **autoimmune** diseases)

IT Glycoproteins, specific or class
 RL: BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (gp39, human chondrocyte protein; autoantigenic **peptides** for inducing tolerance for immunotherapy of **autoimmune** diseases)

IT Immune tolerance
 (specific T cell; autoantigenic **peptides** for inducing tolerance for immunotherapy of **autoimmune** diseases)

IT T cell (lymphocyte)
 (specific tolerance; autoantigenic **peptides** for inducing tolerance for immunotherapy of **autoimmune** diseases)

IT 9001-06-3, Chitotriosidase
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (autoantigenic **peptides** for inducing tolerance for immunotherapy of **autoimmune** diseases)

IT 178274-47-0 256450-45-0 256450-46-1 256450-47-2
 RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (autoantigenic **peptides** for inducing tolerance for immunotherapy of **autoimmune** diseases)

IT 148740-87-8
 RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (pos. control; autoantigenic **peptides** for inducing tolerance for immunotherapy of **autoimmune** diseases)

L4 ANSWER 3 OF 10 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 2000:84638 CAPLUS
 DOCUMENT NUMBER: 132:121462
 TITLE: Use of human cartilage (HC) gp-39 in immune diseases
 INVENTOR(S): Miltenburg, Andreas Martinus Maria; Boots, Anna Maria Helena
 PATENT ASSIGNEE(S): Akzo Nobel N.V., Neth.
 SOURCE: PCT Int. Appl., 29 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000004917	A2	20000203	WO 1999-EP5331	19990719
WO 2000004917	A3	20000720		
W: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
AU 9952875	A1	20000214	AU 1999-52875	19990719
EP 1100526	A2	20010523	EP 1999-938340	19990719
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				

PRIORITY APPLN. INFO.: EP 1998-202471 A 19980723
 WO 1999-EP5331 W 19990719

AB The present invention relates to the use of HC gp-39 to prevent (auto)immune disease or inflammatory diseases, e.g. rheumatoid arthritis. More specifically, HC gp-39 or fragments thereof can be used to modulate the reactivity of lymphocytes which are reactive to antigens other than HC gp-39 but which are present in the same tissue as where HC gp-39 is being expressed.

IN Miltenburg, Andreas Martinus Maria; Boots, Anna Maria Helena
 ST human cartilage gp39 **autoimmune** inflammatory disease; rheumatoid arthritis gp39 **peptide**

IT Glycoproteins, specific or class
 RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (CMP (cartilage matrix protein), gp-39; human cartilage gp-39 **peptides** for preventing (auto)immune and inflammatory diseases)

IT Immunity
 (disorder; human cartilage gp-39 **peptides** for preventing (auto)immune and inflammatory diseases)

IT Glycoproteins, specific or class
 RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (gp39; human cartilage gp-39 **peptides** for preventing (auto)immune and inflammatory diseases)

IT **Autoimmune** disease
 Inflammation
 Protein sequences
 Rheumatoid arthritis
 (human cartilage gp-39 **peptides** for preventing (auto)immune and inflammatory diseases)

IT Lymphocyte

(modulation; human cartilage gp-39 peptides for preventing (auto)immune and inflammatory diseases)
IT 178274-42-5 178274-43-6 178274-44-7 178274-45-8 178274-46-9
178274-47-0 178274-48-1 178274-49-2
RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(human cartilage gp-39 peptides for preventing (auto)immune and inflammatory diseases)

L4 ANSWER 4 OF 10 MEDLINE MEDLINE DUPLICATE 1

ACCESSION NUMBER: 1999250342 MEDLINE
DOCUMENT NUMBER: 99250342 PubMed ID: 10233745
TITLE: T-cell anergy induced by clonotype-specific antibodies: modulation of an autoreactive human T-cell clone in vitro.
AUTHOR: Steenbakkers P G; Boots A M; Rijnders A W
CORPORATE SOURCE: Department of Immunology, N. V. Organon, Oss, The Netherlands.
SOURCE: IMMUNOLOGY, (1999 Apr) 96 (4) 586-94.
PUB. COUNTRY: Journal code: 0374672. ISSN: 0019-2805.
ENGLAND: United Kingdom
LANGUAGE: Journal; Article; (JOURNAL ARTICLE)
FILE SEGMENT: English
ENTRY MONTH: Priority Journals
ENTRY DATE: 199907
Entered STN: 19990816
Last Updated on STN: 19990816
Entered Medline: 19990730

AB Monoclonal antibodies (mAb) specific for the clonotype of an autoreactive T cell may be useful reagents in the modulation of autoimmune disease. We have previously reported the generation of a set of mAb specific for the clonotypic structure of a human T-cell clone recognizing an epitope of human cartilage gp-39. This glycoprotein was recently identified as a candidate autoantigen in rheumatoid arthritis. Here, we demonstrate for the first time that small amounts of immobilized anticlotype mAb can induce anergy in the autoreactive clone. Following the anergic stimulus, T cells failed to proliferate upon restimulation as a result of a lack of interleukin-2 (IL-2) gene transcription. In addition, a diminished interferon-gamma (IFN-gamma) production was found. Our data indicate that anergy was not a result of T-cell receptor (TCR) downmodulation or the absence of free TCR. The anergic state was induced independent of costimulation or the presence of IL-2 and no protein synthesis was required for the induction of anergy. Anticlotype mAb-induced anergy was prevented by cyclosporin A, suggesting that active signalling via the calcium/calmodulin pathway was required for the induction of anergy. In coculture experiments, anergic T cells were found to suppress the response of reactive cells from the same clone. This bystander suppression led to 90% inhibition of peptide-induced proliferation. Together, these findings suggest that mAb to the clonotypic structure of autoreactive T cells may be suitable reagents for the functional inactivation of these T cells in autoimmune diseases.

AU Steenbakkers P G; Boots A M; Rijnders A W
AB Monoclonal antibodies (mAb) specific for the clonotype of an autoreactive T cell may be useful reagents in the modulation of autoimmune disease. We have previously reported the generation of a set of mAb specific for the clonotypic structure of a human. . . . found to suppress the response of reactive cells from the same clone. This bystander suppression led to 90% inhibition of peptide-induced proliferation. Together, these findings suggest that mAb to the clonotypic structure of autoreactive T cells may be suitable reagents for the functional inactivation of these T cells in autoimmune diseases.

L4 ANSWER 5 OF 10 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 2

ACCESSION NUMBER: 1998:785564 CAPLUS
DOCUMENT NUMBER: 130:37290
TITLE: Proteins and novel peptides derived from autoantigen for use in immunotherapy of autoimmune diseases
INVENTOR(S): Boots, Anna Maria Helena; Verheijden, Gijbertus Franciscus Maria; Bos, Ebo Sybren
PATENT ASSIGNEE(S): Akzo Nobel N.V., Neth.
SOURCE: U.S., 19 pp., Cont.-in-part of U.S. 5,736,507.
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 3
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5843449	A	19981201	US 1996-634493	19960418
US 5736507	A	19980407	US 1996-619645	19960325
CA 2251584	AA	19971030	CA 1997-2251584	19970415
WO 9740149	A1	19971030	WO 1997-EP1903	19970415
W:	AM, AU, BB, BG, BR, CA, CN, CZ, EE, GE, HU, IS, JP, KG, KP, KR, LK, LR, LT, LV, MD, MG, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TR, TT, UA, US, UZ, VN, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CP, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
AU 9723869	A1	19971112	AU 1997-23869	19970415
AU 724547	B2	20000928		
EP 904369	A1	19990331	EP 1997-919370	19970415
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI			
CN 1216582	A	19990512	CN 1997-193881	19970415
BR 9708714	A	19990803	BR 1997-8714	19970415
JP 2000059265	T2	20000725	JP 1997-537477	19970415
KR 2000005476	A	20000125	KR 1998-8253	19981015
NO 9804835	A	19981216	NO 1998-4835	19981016
PRIORITY APPLN. INFO.:			US 1996-619645	A2 19960325
			EP 1994-203128	A 19941027
			EP 1995-200886	A 19950407
			WO 1995-EP4201	W 19951025
			US 1996-634493	A 19960418
			WO 1997-EP1903	W 19970415

AB The present invention relates to novel peptides derived from the autoantigen HC gp-39, said peptides comprising at least one of the amino acid sequences FGSRPTILAS (SEQ ID No. 1), FTLASSETG (SEQ ID No. 2), YDDQESVKS (SEQ ID No. 3) and FSKIASNTQ (SEQ ID No. 4). The peptides resemble MHC Class II restricted T-cell epitopes present on the autoantigen HC gp-39 in articular cartilage. HC gp-39, proteins comprising an amino acid sequence which exhibits at least 50% homol. with the amino acid sequence YKLVCCYTSWSQYREGDSCFPDADRFLCTHIYSFANISND (SEQ

ID No: 10) and said **peptides** can be used in antigen-specific treatment of articular cartilage destruction in **autoimmune** diseases in mammals to induce systemic tolerance of the immune system. The autoantigen HC gp-39, proteins comprising an amino acid sequence which exhibits at least 50% homol. with the amino acid sequence YKLVCYTSWSQYREGDGSCFPDALDRFLCTHIIYSFANISND (SEQ ID NO: 10) and said **peptides** are also suitable to induce arthritis in animals, preferably mice. The invention furthermore relates to pharmaceutical compns. comprising said autoantigen and/or said **peptides**, a diagnostic method for the detection of autoreactive T cells in a test sample and test kits to be used in said method. For treating T cell-mediated cartilage destruction disease (e.g. arthritis or rheumatoid arthritis), the T cell-specific tolerance-inducing **peptide** or protein can also be selected from the group consisting of pig heparin binding 38 kDa protein, bovine 39 kDa whey protein, murine breast regressing 39 kDa protein (brp39), human oviduct-specific glycoprotein, murine oviduct-specific glycoprotein, hamster oviduct-specific glycoprotein, bovine oviduct-specific glycoprotein, human chitotriosidase precursor and their fragments.

REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

- TI Proteins and novel **peptides** derived from autoantigen for use in immunotherapy of **autoimmune** diseases
- IN Boots, Anna Maria Helena; Verheijden, Gijsbertus Franciscus Maria; Bos, Ebo Sybren
- AB The present invention relates to novel **peptides** derived from the autoantigen HC gp-39, said **peptides** comprising at least one of the amino acid sequences FGSRPTILAS (SEQ ID No. 1), FTLASSETG (SEQ ID No. 2), YDQESVKS (SEQ ID No. 3) and PSKIASNTQ (SEQ ID No. 4). The **peptides** resemble MHC Class II restricted T-cell epitopes present on the autoantigen HC gp-39 in articular cartilage. HC gp-39, proteins comprising an amino acid sequence which exhibits at least 50% homol. with the amino acid sequence YKLVCYTSWSQYREGDGSCFPDALDRFLCTHIIYSFANISND (SEQ ID No: 10) and said **peptides** can be used in antigen-specific treatment of articular cartilage destruction in **autoimmune** diseases in mammals to induce systemic tolerance of the immune system. The autoantigen HC gp-39, proteins comprising an amino acid sequence which exhibits at least 50% homol. with the amino acid sequence YKLVCYTSWSQYREGDGSCFPDALDRFLCTHIIYSFANISND (SEQ ID No: 10) and said **peptides** are also suitable to induce arthritis in animals, preferably mice. The invention furthermore relates to pharmaceutical compns. comprising said autoantigen and/or said **peptides**, a diagnostic method for the detection of autoreactive T cells in a test sample and test kits to be used in said method. For treating T cell-mediated cartilage destruction disease (e.g. arthritis or rheumatoid arthritis), the T cell-specific tolerance-inducing **peptide** or protein can also be selected from the group consisting of pig heparin binding 38 kDa protein, bovine 39 kDa whey protein, murine breast regressing 39 kDa protein (brp39), human oviduct-specific glycoprotein, murine oviduct-specific glycoprotein, hamster oviduct-specific glycoprotein, bovine oviduct-specific glycoprotein, human chitotriosidase precursor and their fragments.
- ST autoantigen gp39 **autoimmune** cartilage destruction disease; rheumatoid arthritis autoantigen HC gp39 **peptide**
- IT Proteins, specific or class
RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(38,000-mol.-wt.; **peptides** derived from autoantigen HC gp39 for inducing T cell-specific tolerance and for treating **autoimmune** disease such as cartilage destruction diseases and arthritis or rheumatoid arthritis)
- IT Proteins, specific or class
RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(39,000-mol.-wt.; **peptides** derived from autoantigen HC gp39 for inducing T cell-specific tolerance and for treating **autoimmune** disease such as cartilage destruction diseases and arthritis or rheumatoid arthritis)
- IT Histocompatibility antigens
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
(MHC (major histocompatibility complex), class II; **peptides** derived from autoantigen HC gp39 for inducing T cell-specific tolerance and for treating **autoimmune** disease such as cartilage destruction diseases and arthritis or rheumatoid arthritis)
- IT Epitopes
(T cell; **peptides** derived from autoantigen HC gp39 for inducing T cell-specific tolerance and for treating **autoimmune** disease such as cartilage destruction diseases and arthritis or rheumatoid arthritis)
- IT Antigens
RL: ADV (Adverse effect, including toxicity); ARU (Analytical role, unclassified); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study)
(autoantigens, HC gp39; **peptides** derived from autoantigen HC gp39 for inducing T cell-specific tolerance and for treating **autoimmune** disease such as cartilage destruction diseases and arthritis or rheumatoid arthritis)
- IT Proteins, specific or class
RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(breast regressing 39 kDa proteins; **peptides** derived from autoantigen HC gp39 for inducing T cell-specific tolerance and for treating **autoimmune** disease such as cartilage destruction diseases and arthritis or rheumatoid arthritis)
- IT Cartilage
Cartilage
(degeneration; **peptides** derived from autoantigen HC gp39 for inducing T cell-specific tolerance and for treating **autoimmune** disease such as cartilage destruction diseases and arthritis or rheumatoid arthritis)
- IT T cell (lymphocyte)
(epitope; **peptides** derived from autoantigen HC gp39 for inducing T cell-specific tolerance and for treating **autoimmune** disease such as cartilage destruction diseases and arthritis or rheumatoid arthritis)
- IT Oviduct
(glycoprotein specific to; **peptides** derived from autoantigen HC gp39 for inducing T cell-specific tolerance and for treating **autoimmune** disease such as cartilage destruction diseases and arthritis or rheumatoid arthritis)
- IT Proteins, specific or class
RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL

(Biological study); USES (Uses)
 (heparin-binding, 38 kDa; **peptides** derived from autoantigen HC gp39 for inducing T cell-specific tolerance and for treating autoimmune disease such as cartilage destruction diseases and arthritis or rheumatoid arthritis)

IT Drug delivery systems
 (injections; **peptides** derived from autoantigen HC gp39 for inducing T cell-specific tolerance and for treating autoimmune disease such as cartilage destruction diseases and arthritis or rheumatoid arthritis)

IT Drug delivery systems
 (nasal, intra-; **peptides** derived from autoantigen HC gp39 for inducing T cell-specific tolerance and for treating autoimmune disease such as cartilage destruction diseases and arthritis or rheumatoid arthritis)

IT Drug delivery systems
 (oral; **peptides** derived from autoantigen HC gp39 for inducing T cell-specific tolerance and for treating autoimmune disease such as cartilage destruction diseases and arthritis or rheumatoid arthritis)

IT Glycoproteins, general, biological studies
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (oviduct-specific; **peptides** derived from autoantigen HC gp39 for inducing T cell-specific tolerance and for treating autoimmune disease such as cartilage destruction diseases and arthritis or rheumatoid arthritis)

IT Animal
 Arthritis
 Autoimmune disease
 Cattle
 Hamster
 Immune tolerance
 Mammal (Mammalia)
 Mouse
 Protein sequences
 Rheumatoid arthritis
 Rodent
 (**peptides** derived from autoantigen HC gp39 for inducing T cell-specific tolerance and for treating autoimmune disease such as cartilage destruction diseases and arthritis or rheumatoid arthritis)

IT Proteins, specific or class
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (secretory, YM-1 precursor; **peptides** derived from autoantigen HC gp39 for inducing T cell-specific tolerance and for treating autoimmune disease such as cartilage destruction diseases and arthritis or rheumatoid arthritis)

IT Proteins, specific or class
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (whey, 39 kDa; **peptides** derived from autoantigen HC gp39 for inducing T cell-specific tolerance and for treating autoimmune disease such as cartilage destruction diseases and arthritis or rheumatoid arthritis)

IT 198841-51-9
 RL: ADV (Adverse effect, including toxicity); ARU (Analytical role, unclassified); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study)
 (**peptides** derived from autoantigen HC gp39 for inducing T cell-specific tolerance and for treating autoimmune disease such as cartilage destruction diseases and arthritis or rheumatoid arthritis)

IT 172253-32-6
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (**peptides** derived from autoantigen HC gp39 for inducing T cell-specific tolerance and for treating autoimmune disease such as cartilage destruction diseases and arthritis or rheumatoid arthritis)

IT 148740-87-8 178274-42-5 178274-43-6 178274-44-7 178274-45-8 178274-46-9 178274-47-0 178274-48-1 178274-49-2
 RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (**peptides** derived from autoantigen HC gp39 for inducing T cell-specific tolerance and for treating autoimmune disease such as cartilage destruction diseases and arthritis or rheumatoid arthritis)

L4 ANSWER 6 OF 10 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
 ACCESSION NUMBER: 2002:106274 BIOSIS
 DOCUMENT NUMBER: PREV200200106274
 TITLE: **Peptides** derived from autoantigen for use in immunotherapy of autoimmune diseases.
 AUTHOR(S): Boots, A. M. H.; Verheijden, G. F. M.
 CORPORATE SOURCE: Verlengde Torenstraat Netherlands
 ASSIGNEE: AKZO NOBEL N.V.
 PATENT INFORMATION: US 5736507 April 7, 1998
 SOURCE: Official Gazette of the United States Patent and Trademark Office Patents, (April 7, 1998) Vol. 1209, No. 1, pp. 492.
 ISSN: 0098-1133.
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 TI **Peptides** derived from autoantigen for use in immunotherapy of autoimmune diseases.
 AU Boots, A. M. H.; Verheijden, G. F. M.
 IT Sequence Data
 AMINO ACID SEQUENCE
 IT Miscellaneous Descriptors
 CHEMICAL FORMULA; HUMAN CARTILAGE PEPTIDE; IMMUNOLOGIC AGENT; PHARMACEUTICALS

L4 ANSWER 7 OF 10 CAPLUS COPYRIGHT 2002 ACS
 ACCESSION NUMBER: 1997:717938 CAPLUS
 DOCUMENT NUMBER: 128:2901
 TITLE: Novel **peptides** suitable for use in antigen specific immunosuppressive therapy
 INVENTOR(S): Boots, Anna Maria Helena; Verheijden, Gijsbertus Franciscus Maria
 PATENT ASSIGNEE(S): Akzo Nobel N.V., Neth.; Boots, Anna Maria Helena; Verheijden, Gijsbertus Franciscus Maria
 SOURCE: PCT Int. Appl., 82 pp.

DOCUMENT TYPE: CODEN: PIXXD2
LANGUAGE: Patent
FAMILY ACC. NUM. COUNT: 1 English
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9740068	A1	19971030	WO 1997-EP2051	19970422
W: AU, BR, CA, CN, CZ, HU, JP, KR, MX, NO, NZ, PL, RU, SG, TR, US				
RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
ZA 9703071	A	19980805	ZA 1997-3071	19970410
CA 2251680	AA	19971030	CA 1997-2251680	19970422
AU 9727685	A1	19971112	AU 1997-27685	19970422
AU 719481	B2	20000511		
CN 1216551	A	19990512	CN 1997-194031	19970422
EP 920451	A1	19990609	EP 1997-921707	19970422
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
BR 9708744	A	19990803	BR 1997-8744	19970422
JP 2000510106	T2	20000808	JP 1997-537744	19970422
KR 2000010561	A	20000215	KR 1998-708415	19981021
NO 9804937	A	19981201	NO 1998-4937	19981023
US 6184204	B1	20010206	US 1998-171705	19981023
PRIORITY APPLN. INFO.: EP 1996-201106 A 19960424 WO 1997-EP2051 W 19970422				

AB The invention relates to **peptides** consisting of 16 to 55 amino acid residues, and useful in diagnosis for detecting activated autoreactive T cells in the individual. The **peptides** are also useful in the treatment of **autoimmune** disease, e.g. T-cell mediated destruction of articular cartilage. Administration of pharmaceutical compns. based on these **peptides** can be used to induce systemic immunol. tolerance to the autoantigens under attack of the autoreactive T-cells.

TI Novel **peptides** suitable for use in antigen specific immunosuppressive therapy

IN Boots, Anna Maria Helena; Verheijden, Gijsbertus Franciscus Maria

AB The invention relates to **peptides** consisting of 16 to 55 amino acid residues, and useful in diagnosis for detecting activated autoreactive T cells in the individual. The **peptides** are also useful in the treatment of **autoimmune** disease, e.g. T-cell mediated destruction of articular cartilage. Administration of pharmaceutical compns. based on these **peptides** can be used to induce systemic immunol. tolerance to the autoantigens under attack of the autoreactive T-cells.

ST antigen immunosuppressant autoreactive T cell tolerance; **autoimmune** disease articular cartilage immunosuppressive **peptide**

IT Cartilage
(articular, destruction; immunosuppressive antigen **peptides** for inhibiting autoreactive T cell and for treating and diagnosing **autoimmune** disease)

IT Antigens
RL: BSU (Biological study, unclassified); BIOL (Biological study) (autoantigens; immunosuppressive antigen **peptides** for inhibiting autoreactive T cell and for treating and diagnosing **autoimmune** disease)

IT T cell (lymphocyte)
(autoreactive; immunosuppressive antigen **peptides** for inhibiting autoreactive T cell and for treating and diagnosing **autoimmune** disease)

IT **Autoimmune** disease
Immune tolerance
Immunosuppressants
Protein sequences
(immunosuppressive antigen **peptides** for inhibiting autoreactive T cell and for treating and diagnosing **autoimmune** disease)

IT Antigens
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses) (immunosuppressive antigen **peptides** for inhibiting autoreactive T cell and for treating and diagnosing **autoimmune** disease)

IT Mononuclear cell (leukocyte)
(peripheral blood; immunosuppressive antigen **peptides** for inhibiting autoreactive T cell and for treating and diagnosing **autoimmune** disease)

IT Blood
(peripheral; immunosuppressive antigen **peptides** for inhibiting autoreactive T cell and for treating and diagnosing **autoimmune** disease)

IT 198880-51-2 198880-52-3 198880-53-4 198880-54-5 198880-55-6
198880-56-7 198880-57-8 198880-58-9 198880-59-0 198880-60-3
198880-61-4 198880-62-5 198880-63-6 198880-64-7 198880-65-8
198880-67-0 198880-69-2 198880-70-5 198880-71-6 198880-72-7
198880-73-8 198880-74-9 198880-75-0 198880-76-1 198880-77-2
198880-78-3 198880-79-4 198880-80-7 198880-81-8 198880-82-9
198880-83-0 198880-84-1 198880-85-2 198880-86-3 198880-87-4
198880-88-5 198880-89-6 198880-90-9 198880-91-0 198880-92-1
198880-93-2 198880-94-3 198880-95-4 198880-96-5

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(immunosuppressive antigen **peptides** for inhibiting autoreactive T cell and for treating and diagnosing **autoimmune** disease)

L4 ANSWER 8 OF 10 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1997:151549 CAPLUS

DOCUMENT NUMBER: 126:152794

TITLE: **Peptides** for use in treatment of T-cell mediated cartilage destruction in **autoimmune** diseases

INVENTOR(S): Verheijden, Gijsbertus Francisc; Boots, Anna Maria Helena

PATENT ASSIGNEE(S): Akzo Nobel N.V., Neth.; Verheijden, Gijsbertus Franciscus Maria; Boots, Anna Maria Helena

SOURCE: PCT Int. Appl., 22 pp.
CODEN: PIXXD2

DOCUMENT TYPE: Patent
LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9700270	A1	19970103	WO 1996-EP2605	19960617
W: AL, AM, AU, BB, BG, BR, CA, CN, CZ, EE, GE, HU, IS, JP, KG, KP, KR, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
CA 2221981	AA	19970103	CA 1996-2221981	19960617
AU 9662246	A1	19970115	AU 1996-62246	19960617
AU 704942	B2	19990506		
EP 833842	A1	19980408	EP 1996-920822	19960617
EP 833842	B1	19990929		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
CN 1188483	A	19980722	CN 1996-194869	19960617
BR 9608626	A	19990615	BR 1996-8626	19960617
JP 11507919	T2	19990713	JP 1996-502649	19960617
AT 185151	E	19991015	AT 1996-920822	19960617
ES 2139363	T3	20000201	ES 1996-920822	19960617
ZA 9605163	A	19970123	ZA 1996-5163	19960618
NO 9705968	A	19971218	NO 1997-5968	19971218
PRIORITY APPLN. INFO.: EP 1995-201656 A 19950619 WO 1996-EP2605 W 19960617				
AB	The invention relates to the use of novel peptides in a peptide-induced tolerance therapy for the induction of tolerance to autoaggressive T cells assocd. with T-cell mediated articular cartilage destruction in autoimmune diseases, more specifically arthritis. The invention furthermore embraces pharmaceutical compns. comprising said peptides and a diagnostic method for the detection of autoreactive T cells in a test sample, said T cells being assocd. with T-cell mediated articular cartilage destruction in autoimmune diseases and test kits to be used in said method.			
TI	Peptides for use in treatment of T-cell mediated cartilage destruction in autoimmune diseases			
IN	Verheijden, Gijbertus Francisc; Boots, Anna Maria Helena			
AB	The invention relates to the use of novel peptides in a peptide-induced tolerance therapy for the induction of tolerance to autoaggressive T cells assocd. with T-cell mediated articular cartilage destruction in autoimmune diseases, more specifically arthritis. The invention furthermore embraces pharmaceutical compns. comprising said peptides and a diagnostic method for the detection of autoreactive T cells in a test sample, said T cells being assocd. with T-cell mediated articular cartilage destruction in autoimmune diseases and test kits to be used in said method.			
ST	autoimmune disease cartilage destruction peptide therapy			
IT	Autoimmune disease Cartilage Rheumatoid arthritis T cell (lymphocyte) (peptides for use in treatment of T-cell mediated cartilage destruction in autoimmune diseases)			
IT	186827-57-6 186827-58-7 186827-59-8 186827-60-1 RI: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (peptides for use in treatment of T-cell mediated cartilage destruction in autoimmune diseases)			
L4	ANSWER 9 OF 10 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.			
ACCESSION NUMBER:	1998037624 EMBASE			
TITLE:	Selection of self-reactive peptides within human aggrecan by use of a HLA-DRB1*0401 peptide binding motif.			
AUTHOR:	Boots A.M.H.; Verheijden G.F.M.; Schoningh R.; Van Staveren C.J.; Bos E.; Elewaut D.; De Keyser F.; Veys E.; Joosten I.; Rijnders A.W.M.			
CORPORATE SOURCE:	Dr. A.M.H. Boots, Department of Immunology, RH165, NV Organon, P.O. Box 20, 5340 BH Oss, Netherlands			
SOURCE:	Journal of Autoimmunity, (1997) 10/6 (569-578). Refs: 40 ISSN: 0896-8411 CODEN: JOAUPE			
COUNTRY:	United Kingdom			
DOCUMENT TYPE:	Journal; Article			
FILE SEGMENT:	026 Immunology, Serology and Transplantation			
LANGUAGE:	English			
SUMMARY LANGUAGE:	English			
AB	The pathogenesis of joint destruction in rheumatoid arthritis remains ill-defined. Joint destruction is thought to be the result of tissue damage mediated by T cells. The mere presence of articular cartilage appears responsible for sustaining chronic synovitis and thereby forwards a role for cartilage-responsive T cells in RA. Taking advantage of the positive DRB1*0401 association with RA susceptibility, we reasoned that T-cell recognition of autoantigens in RA would be restricted by DRB1*0401-encoded molecules. A DR4 (B1*0401) peptide binding motif was used for the identification of putative T-cell epitopes within human aggrecan, a candidate autoantigen. Thirteen peptides were synthesized and tested for binding DRB1*0401 or 0404-encoded molecules. Selected binders were tested for induction of proliferative responses in peripheral blood mononuclear cells from donors carrying the DR4 or DR1 specificity. Both healthy and RA donors responded to human aggrecan-derived peptides , thereby identifying these sequences as T-cell epitopes. Interestingly, responses to aggrecan-derived epitopes were significantly decreased in RA patients compared to controls. This was not due to an overall hyporesponsiveness of RA patients since responses to a recall antigen or mitogen did not differ from controls. The data suggest that in RA, aggrecan-specific T cells may exist in a different stage of activation or may have left the periphery to home to the joint.			
TI	Selection of self-reactive peptides within human aggrecan by use of a HLA-DRB1*0401 peptide binding motif.			
AU	Boots A.M.H.; Verheijden G.F.M.; Schoningh R.; Van Staveren C.J.; Bos E.; Elewaut D.; De Keyser F.; Veys E.; Joosten I.; Rijnders A.W.M.			
AB	. . . RA susceptibility, we reasoned that T-cell recognition of autoantigens in RA would be restricted by DRB1*0401-encoded molecules. A DR4 (B1*0401) peptide binding motif was used for the identification of putative T-cell epitopes within human aggrecan, a candidate autoantigen. Thirteen peptides were synthesized and tested for binding DRB1*0401 or 0404-encoded molecules. Selected binders were tested for induction of proliferative responses in. . . blood			

mononuclear cells from donors carrying the DR4 or DR1 specificity. Both healthy and RA donors responded to human aggrecan-derived **peptides**, thereby identifying these sequences as T-cell epitopes. Interestingly, responses to aggrecan-derived epitopes were significantly decreased in RA patients compared to . . .

CT Medical Descriptors:

***peptide analysis**
 autoimmunity
 t lymphocyte activation
 pathogenesis
 tissue injury
 molecular recognition
 mononuclear cell
 antigen specificity
 antigen presentation
 cell proliferation
 human
 human cell
 article
 priority journal
 *aggrecan
 *HLA DR antigen

L4 ANSWER 10 OF 10 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1996:397369 CAPLUS

DOCUMENT NUMBER: 125:49310

TITLE: Novel **peptides** derived from the articular cartilage autoantigen HC gp-39 for use in immunotherapy of **autoimmune** diseases

INVENTOR(S): Boots, Anna Maria Helena; Verheijden, Gijsbertus Franciscus Maria

PATENT ASSIGNEE(S): Akzo Nobel N.V., Neth.

SOURCE: PCT Int. Appl., 37 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9613517	A1	19960509	WO 1995-EP4201	19951025
W: AU, BR, CA, CN, FI, HU, JP, KR, MX, NO, NZ, PL, RU, US				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
IL 115744	A1	20000716	IL 1995-115744	19951024
AU 9539252	A1	19960523	AU 1995-39252	19951025
AU 696827	B2	19980917		
EP 733065	A1	19960925	EP 1995-937008	19951025
EP 733065	B1	19990317		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
HU 74847	A2	19970228	HU 1996-1401	19951025
HU 218027	B	20000528		
JP 09507861	T2	19970812	JP 1995-514306	19951025
BR 9506377	A	19970916	BR 1995-6377	19951025
CN 1168677	A	19971224	CN 1995-190944	19951025
AT 177756	E	19990415	AT 1995-937008	19951025
ES 2130672	T3	19990701	ES 1995-937008	19951025
ZA 9509123	A	19970123	ZA 1995-9123	19951027
US 5736507	A	19980407	US 1996-619645	19960325
FI 9602619	A	19960625	FI 1996-2619	19960625
NO 9602695	A	19960626	NO 1996-2695	19960626
PRIORITY APPLN. INFO.:			EP 1994-203128	A 19941027
			EP 1995-200886	A 19950407
			WO 1995-EP4201	W 19951025

AB Novel **peptides** derived from the autoantigen HC gp-39 including at least one of the fragments FGRSFTLAS, FTLASSETC, YDDQESVKS or FSKIASNTQ are described for use in the induction of immune tolerance in the treatment of **autoimmune** disease. The **peptides** resemble MHC Class II restricted T-cell epitopes present on the autoantigen HC gp-39 in articular cartilage. HC gp-39 and these **peptides** can be used in antigen-specific treatment of articular cartilage destruction in **autoimmune** diseases to induce tolerance of the immune system. The autoantigen HC gp-39 and these **peptides** are also suitable to induce arthritis in non-human animals, preferably mice. The invention furthermore relates to pharmaceutical compns. comprising said autoantigen and/or said **peptides**, a diagnostic method for the detection of autoreactive T cells in a test sample and test kits to be used in said method. The use of these **peptides** to induce immune tolerance in mice is demonstrated.

TI Novel **peptides** derived from the articular cartilage autoantigen HC gp-39 for use in immunotherapy of **autoimmune** diseases

IN Boots, Anna Maria Helena; Verheijden, Gijsbertus Franciscus Maria

AB Novel **peptides** derived from the autoantigen HC gp-39 including at least one of the fragments FGRSFTLAS, FTLASSETC, YDDQESVKS or FSKIASNTQ are described for use in the induction of immune tolerance in the treatment of **autoimmune** disease. The **peptides** resemble MHC Class II restricted T-cell epitopes present on the autoantigen HC gp-39 in articular cartilage. HC gp-39 and these **peptides** can be used in antigen-specific treatment of articular cartilage destruction in **autoimmune** diseases to induce tolerance of the immune system. The autoantigen HC gp-39 and these **peptides** are also suitable to induce arthritis in non-human animals, preferably mice. The invention furthermore relates to pharmaceutical compns. comprising said autoantigen and/or said **peptides**, a diagnostic method for the detection of autoreactive T cells in a test sample and test kits to be used in said method. The use of these **peptides** to induce immune tolerance in mice is demonstrated.

ST gp39 **peptide** autoantigen **autoimmune** disease therapy

IT Immune tolerance
 (induction of; novel **peptides** derived from articular cartilage autoantigen HC gp-39 for use in immunotherapy of **autoimmune** diseases)

IT Lymphocyte
 (T-cell, autoreactive, detection of; novel **peptides** derived from articular cartilage autoantigen HC gp-39 for use in immunotherapy of **autoimmune** diseases)

IT Inflammation inhibitors
 (antiarthritics, novel **peptides** derived from articular cartilage autoantigen HC gp-39 for use in immunotherapy of **autoimmune** diseases)

IT Cartilage
 (articular, preventing destruction in **autoimmune** disease of;

novel peptides derived from articular cartilage autoantigen
 HC gp-39 for use in immunotherapy of autoimmune diseases)

IT Antigens
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (auto-, cartilage HC gp-39; novel peptides derived from
 articular cartilage autoantigen HC gp-39 for use in immunotherapy of
 autoimmune diseases)

IT Glycoproteins, specific or class
 RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES
 (Uses)
 (gp39, cartilage autoantigen; novel peptides derived from
 articular cartilage autoantigen HC gp-39 for use in immunotherapy of
 autoimmune diseases)

IT Arthritis
 (rheumatoid, novel peptides derived from articular cartilage
 autoantigen HC gp-39 for use in immunotherapy of autoimmune
 diseases)

IT 178274-42-5D, analogs, derivs. 178274-43-6D, analogs, derivs.
 178274-44-7D, analogs, derivs. 178274-45-8D, analogs, derivs.
 178274-46-9D, analogs, derivs. 178274-47-0D, analogs, derivs.
 178274-48-1D, analogs, derivs. 178274-49-2D, analogs, derivs.
 RL: PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES
 (Uses)
 (novel peptides derived from articular cartilage autoantigen
 HC gp-39 for use in immunotherapy of autoimmune diseases)

=> end

ALL L# QUERIES AND ANSWER SETS ARE DELETED AT LOGOFF

LOGOFF? (Y)/N/HOLD:y

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
31.95	32.16

FULL ESTIMATED COST

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
ENTRY	SESSION
-4.34	-4.34

CA SUBSCRIBER PRICE

STN INTERNATIONAL LOGOFF AT 22:45:43 ON 24 JUN 2002

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:sssptal644axd

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

***** Welcome to STN International *****

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
 NEWS 2 Jan 25 BLAST(R) searching in REGISTRY available in STN on the Web
 NEWS 3 Jan 29 FSTA has been reloaded and moves to weekly updates
 NEWS 4 Feb 01 DKILIT now produced by FIZ Karlsruhe and has a new update
 frequency
 NEWS 5 Feb 19 Access via Tymnet and SprintNet Eliminated Effective 3/31/02
 NEWS 6 Mar 08 Gene Names now available in BIOSIS
 NEWS 7 Mar 22 TOXLIT no longer available
 NEWS 8 Mar 22 TRCTHERMO no longer available
 NEWS 9 Mar 28 US Provisional Priorities searched with P in CA/Caplus
 and USPATFULL
 NEWS 10 Mar 28 LIPINSKI/CALC added for property searching in REGISTRY
 NEWS 11 Apr 02 PAPERCHEM no longer available on STN. Use PAPERCHEM2 instead.
 NEWS 12 Apr 08 "Ask CAS" for self-help around the clock
 NEWS 13 Apr 09 BEILSTEIN: Reload and Implementation of a New Subject Area
 NEWS 14 Apr 09 ZDB will be removed from STN
 NEWS 15 Apr 19 US Patent Applications available in IFICDB, IFIPAT, and IFIUDB
 NEWS 16 Apr 22 Records from IP.com available in CAPLUS, HCAPLUS, and ZCAPLUS
 NEWS 17 Apr 22 BIOSIS Gene Names now available in TOXCENTER
 NEWS 18 Apr 22 Federal Research in Progress (FEDRIP) now available
 NEWS 19 Jun 03 New e-mail delivery for search results now available
 NEWS 20 Jun 10 MEDLINE Reload
 NEWS 21 Jun 10 PCTFULL has been reloaded

NEWS EXPRESS February 1 CURRENT WINDOWS VERSION IS V6.0d,
 CURRENT MACINTOSH VERSION IS V6.0a(ENG) AND V6.0Ja(JP),
 AND CURRENT DISCOVER FILE IS DATED 05 FEBRUARY 2002

NEWS HOURS STN Operating Hours Plus Help Desk Availability
 NEWS INTER General Internet Information
 NEWS LOGIN Welcome Banner and News Items
 NEWS PHONE Direct Dial and Telecommunication Network Access to STN
 NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that
 specific topic.

All use of STN is subject to the provisions of the STN Customer
 agreement. Please note that this agreement limits use to scientific
 research. Use for software development or design or implementation
 of commercial gateways or other similar uses is prohibited and may
 result in loss of user privileges and other penalties.

***** STN Columbus *****

FILE 'HOME' ENTERED AT 09:46:22 ON 25 JUN 2002

=> file medline caplus embase biosis

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
0.21	0.21

FULL ESTIMATED COST

FILE 'MEDLINE' ENTERED AT 09:46:36 ON 25 JUN 2002

FILE 'CAPLUS' ENTERED AT 09:46:36 ON 25 JUN 2002

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'EMBASE' ENTERED AT 09:46:36 ON 25 JUN 2002
COPYRIGHT (C) 2002 Elsevier Science B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 09:46:36 ON 25 JUN 2002
COPYRIGHT (C) 2002 BIOLOGICAL ABSTRACTS INC.(R)

=> s FTLASAETT or HSFTLASAETTVG
L1 0 FTLASAETT OR HSFTLASAETTVG

=> s ykl
L2 211 YKL

=> s l2 and chondrocyte?
L3 69 L2 AND CHONDROCYTE?

=> s l3 and (autoimmun? or RA or arthritis)
L4 42 L3 AND (AUTOIMMUN? OR RA OR ARTHRITIS)

=> dup rem l4
DUUP IS NOT A RECOGNIZED COMMAND
The previous command name entered was not recognized by the system.
For a list of commands available to you in the current file, enter
"HELP COMMANDS" at an arrow prompt (=>).

=> dup rem l4
PROCESSING COMPLETED FOR L4
L5 21 DUP REM L4 (21 DUPLICATES REMOVED)

=> dis l5 1-21 ibib abs

L5 ANSWER 1 OF 21 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.
ACCESSION NUMBER: 2001147218 EMBASE
TITLE: Regulation of YKL-40 production by human
articular chondrocytes.
AUTHOR: Johansen J.S.; Olee T.; Price P.A.; Hashimoto S.; Ochs
R.L.; Lotz M.
CORPORATE SOURCE: Dr. J.S. Johansen, Department of Internal Medicine,
Division of Rheumatology 232, Hvidovre Hospital, Kettegaard
alle 30, DK-2650 Hvidovre, Denmark
SOURCE: Arthritis and Rheumatism, (2001) 44/4 (826-837).
Refs: 40
ISSN: 0004-3591 CODEN: ARHEAW
COUNTRY: United States
DOCUMENT TYPE: Journal; Article
FILE SEGMENT: 005 General Pathology and Pathological Anatomy
029 Clinical Biochemistry
031 Arthritis and Rheumatism
033 Orthopedic Surgery
LANGUAGE: English
SUMMARY LANGUAGE: English

AB Objective. YKL-40 (human cartilage glycoprotein 39) is one of
the most abundant proteins secreted by cultured chondrocytes.
The objectives of the present study were to identify regulators of
YKL-40 production in cartilage and chondrocytes and to
map the localization of YKL-40 in chondrocytes.
Methods. Human articular chondrocytes and cartilage explants
(obtained from subjects at autopsy, from a tissue bank, and from
osteoarthritis [OA] patients undergoing total joint replacement surgery)
were stimulated with cytokines, growth factors, and other agents.
YKL-40 expression was analyzed by Northern blot and polymerase
chain reaction. YKL-40 secretion into the media was determined
by enzyme-linked immunosorbent assay. Results. YKL-40 production
increased to very high levels during the early phase of
chondrocyte monolayer culture and in normal cartilage explant
cultures as a response to tissue injury. Spontaneous YKL-40
release was higher in OA than in normal cartilage explant cultures. In
chondrocyte monolayer cultures, interleukin-1.beta. (IL-1.beta.)
and transforming growth factor .beta. (TGF.beta.) decreased the levels of
secreted YKL-40, and this was associated with a reduction in
YKL-40 messenger RNA levels. IL-1.beta., but not TGF.beta.,
reduced YKL-40 production in cartilage explant cultures. Media
from explants treated with cycloheximide had no detectable YKL
-40, suggesting that the released protein was newly synthesized.
Immunofluorescence microscopy showed YKL-40 staining in the
Golgi system of the chondrocytes, but YKL-40 could not
be detected in the extracellular matrix. Conclusion. The spontaneous
increase in the production of YKL-40 in the early phase of
culture appears to represent a cellular response to changes in the
extracellular matrix environment. This, coupled with the profound
suppressive effects of IL-1.beta. and TGF.beta. on YKL-40
production, identifies a novel regulatory pattern for this major
chondrocyte-derived protein.

L5 ANSWER 2 OF 21 MEDLINE MEDLINE DUPLICATE 1
ACCESSION NUMBER: 2001682113 MEDLINE
DOCUMENT NUMBER: 21584364 PubMed ID: 11727845
TITLE: Serum YKL-40 concentrations in patients with
early rheumatoid arthritis: relation to joint
destruction.
AUTHOR: Johansen J S; Kirwan J R; Price P A; Sharif M
CORPORATE SOURCE: Department of Rheumatology, Hvidovre Hospital, University
of Copenhagen, Denmark.. julia.johansen@post3.tele.dk
SOURCE: SCANDINAVIAN JOURNAL OF RHEUMATOLOGY, (2001) 30 (5)
297-304.
Journal code: 0321213. ISSN: 0300-9742.
PUB. COUNTRY: Norway
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 200112
ENTRY DATE: Entered STN: 20011203
Last Updated on STN: 20020123
Entered Medline: 20011211

AB OBJECTIVE: YKL-40 is a secretory glycoprotein of
chondrocytes, synovial cells, macrophages, and neutrophils. The
aims were to determine serum YKL-40 in patients with early
rheumatoid arthritis (RA) and seek associations with
early joint erosions. METHODS: YKL-40 was measured by ELISA in
serum samples collected every three month for 36 months from patients with

early RA. The patients were treated with DMARDs and some were allocated to additional prednisolone. RESULTS: Serum YKL-40 was higher in RA patients compared with controls (98 vs. 42 microg/l, $p < 0.001$). The mean serum YKL-40 during the study correlated with the progression in Larsen score (Pearson's test: $p = 0.004$). Patients with a persistently high serum YKL-40 had larger progression in Larsen score compared with patients with normal serum YKL-40 (median progression: 7 vs. 0, $p = 0.003$). CONCLUSION: These data suggest that elevated serum YKL-40 is related to progression in joint destruction in early RA patients.

LS ANSWER 3 OF 21 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
 ACCESSION NUMBER: 2001:562025 BIOSIS
 DOCUMENT NUMBER: PREV200100562025
 TITLE: YKL-40 gene expression is inhibited by cyclosporin A in human osteoblast-like cell line MG63.
 AUTHOR(S): Musacchio, E. (1); Valvasone, C. (1); Pozzuoli, A. (1); Priante, G. (1); Punzi, L. (1); Netto, F. S. (1); Sartori, L. (1)
 CORPORATE SOURCE: (1) Department of Medical and Surgical Sciences, University of Padova, Padova Italy
 SOURCE: Journal of Bone and Mineral Research, (September, 2001) Vol. 16, No. Suppl. 1, pp. S261. print.
 Meeting Info.: Twenty-Third Annual Meeting of the American Society for Bone and Mineral Research Phoenix, Arizona, USA October 12-16, 2001
 ISSN: 0884-0431.
 DOCUMENT TYPE: Conference
 LANGUAGE: English
 SUMMARY LANGUAGE: English

LS ANSWER 4 OF 21 MEDLINE DUPLICATE 2
 ACCESSION NUMBER: 2001325116 MEDLINE
 DOCUMENT NUMBER: 21198242 PubMed ID: 11300743
 TITLE: Studies on YKL-40 in knee joints of patients with rheumatoid arthritis and osteoarthritis.
 AUTHOR: Involvement of YKL-40 in the joint pathology.
 Volck B; Johansen J S; Stoltenberg M; Garbarsch C; Price P A; Ostergaard M; Ostergaard K; Lovgreen-Nielsen P; Sonne-Holm S; Lorenzen I
 CORPORATE SOURCE: Department of Rheumatology, Hvidovre Hospital, University of Copenhagen, Denmark.. b.volck@dadlnet.dk
 SOURCE: OSTEOARTHRITIS AND CARTILAGE, (2001 Apr) 9 (3) 203-14.
 Journal code: 9305697. ISSN: 1063-4584.
 PUB. COUNTRY: England: United Kingdom
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 200106
 ENTRY DATE: Entered STN: 20010611
 Last Updated on STN: 20010821
 Entered Medline: 20010607

AB OBJECTIVE: The presence of YKL-40 (human cartilage glycoprotein 39) in synovium, cartilage and synovial fluid (SF) from knee joints of patients with rheumatoid arthritis and osteoarthritis (OA) were related to histopathological changes in synovium and cartilage and to serum YKL-40 and other biochemical markers. METHODS: The localization of YKL-40 in synovium and cartilage was determined by immunohistochemistry. Synovial inflammation was estimated histologically and by magnetic resonance imaging (MRI). Biochemical markers of inflammation, neutrophil activation and cartilage metabolism were analysed. YKL-40 concentrations in serum and SF were determined by RIA and ELISA. RESULTS: In the synovium YKL-40 positive cells were found in lining and stromal cells (macrophages) and the number of YKL-40 positive cells was related to the degree of synovitis. In arthritic cartilage, YKL-40 was located to chondrocytes. YKL-40 levels in SF were higher in RA patients with moderate/severe or none/slight synovitis of the knee joint compared to OA patients with moderate/severe or none/slight synovitis. SF YKL-40 correlated with the synovial membrane and the joint effusion volumes determined by magnetic resonance imaging (MRI) and with other biochemical markers of intercellular matrix metabolism. SF YKL-40 was higher than serum YKL-40, and a relationship existed between the YKL-40 levels in SF and serum. Intraarticular glucocorticoid injection was followed by clinical remission and a decrease in serum YKL-40, which increased again at clinical relapse. CONCLUSIONS: YKL-40 in SF is derived from cells in the inflamed synovium, chondrocytes and SF neutrophils. Joint derived YKL-40 influences serum YKL-40. YKL-40 may be involved in the pathophysiology of the arthritic processes and reflect local disease activity.
 Copyright 2001 OsteoArthritis Research Society International.

LS ANSWER 5 OF 21 MEDLINE DUPLICATE 3
 ACCESSION NUMBER: 2001474387 MEDLINE
 DOCUMENT NUMBER: 21408823 PubMed ID: 11518039
 TITLE: Increased level of YKL-40 in sera from patients with early rheumatoid arthritis: a new marker for disease activity.
 AUTHOR: Peltomaa R; Paimela L; Harvey S; Helve T; Leirisalo-Repo M
 CORPORATE SOURCE: Department of Medicine, Helsinki University Central Hospital, Finland.. Ritva.Peltomaa@hus.fi
 SOURCE: RHEUMATOLOGY INTERNATIONAL, (2001 Jul) 20 (5) 192-6.
 Journal code: 8206885. ISSN: 0172-8172.
 PUB. COUNTRY: Germany: Germany, Federal Republic of
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 200201
 ENTRY DATE: Entered STN: 20010827
 Last Updated on STN: 20020125
 Entered Medline: 20020108

AB YKL-40 is a newly discovered major secretory protein of human chondrocytes and synoviocytes. We measured serum levels of YKL-40 in 52 patients with early onset rheumatoid arthritis (RA) by enzyme-linked immunosorbent assay (ELISA) during a 2-year prospective follow-up, correlating values with laboratory and clinical variables and radiographic progression. Levels at baseline before antirheumatic therapy were significantly higher in patients than in healthy controls. The levels of YKL-40 correlated with laboratory and clinical markers of disease activity both at baseline and during follow-up. Baseline YKL-40 values correlated with baseline Larsen scores but did not predict radiographic

progression. Baseline and mean YKL-40 values did not differ between fast and slow radiological progressions. Mean YKL-40 levels correlated with the number of swollen joints but were not predictors of radiographic progression. These results suggest that in early RA, serum YKL-40 is an inflammatory marker correlating with disease activity. However, its levels do not predict clinical course or radiographic progression.

L5 ANSWER 6 OF 21 MEDLINE DUPLICATE 4
 ACCESSION NUMBER: 2001124047 MEDLINE
 DOCUMENT NUMBER: 20566402 PubMed ID: 11114282
 TITLE: Recognition of YKL-39, a human cartilage related protein, as a target antigen in patients with rheumatoid arthritis.
 AUTHOR: Sekine T; Masuko-Hongo K; Matsui T; Asahara H; Takigawa M; Nishioka K; Kato T
 CORPORATE SOURCE: Rheumatology, Immunology and Genetics Programme, Institute of Medical Science, St Marianna University, School of Medicine 2-16-1, Sugao, Miyamae-ku, Kawasaki, Kanagawa, 216-8512, Japan.
 SOURCE: ANNALS OF THE RHEUMATIC DISEASES, (2001 Jan) 60 (1) 49-54. Journal code: 0372355. ISSN: 0003-4967.
 PUB. COUNTRY: ENGLAND: United Kingdom
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 200102
 ENTRY DATE: Entered STN: 20010322
 Last Updated on STN: 20010322
 Entered Medline: 20010222

AB OBJECTIVE: To investigate whether autoimmunity to YKL-39, a recently cloned cartilage protein, occurs in patients with rheumatoid arthritis (RA). METHODS: Autoantibody to YKL-39 was assayed by enzyme linked immunosorbent assay (ELISA) and western blotting in serum samples from patients with RA, systemic lupus erythematosus (SLE), and healthy donors, using recombinant YKL-39 protein. This reactivity was compared with that against a YKL-39 homologue, YKL-40 (human cartilage gp-39/chondrex), which has been reported to be an autoantigen in RA. RESULTS: Autoantibody to YKL-39 was detected in seven of 87 patients with RA (8%), but not in serum samples from patients with SLE or healthy donors. YKL-40 reactivity was found in only one of 87 RA serum samples (1%), with no cross reactivity to YKL-39. CONCLUSION: The existence of anti-YKL-39 antibody in a subset of patients with RA is reported here for the first time. Further, it was shown that the immune response to YKL-39 was independent of that to YKL-40. Clarification of the antibody and T cell responses to autoantigens derived from chondrocyte, cartilage, or other joint components may lead to a better understanding of the pathophysiology of joint destruction in patients with RA.

L5 ANSWER 7 OF 21 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
 ACCESSION NUMBER: 2000:316680 BIOSIS
 DOCUMENT NUMBER: PREV200000316680
 TITLE: YKL-40 and graft rejection.
 AUTHOR(S): Fiore, Carmelo E. (1); Pennisi, Pietra (1); Tamborino, Corrado
 CORPORATE SOURCE: (1) Department of Internal Medicine, University of Catania, Catania Italy
 SOURCE: American Journal of Medicine, (June 1, 2000) Vol. 108, No. 8, pp. 688-689. print. ISSN: 0002-9343.
 DOCUMENT TYPE: Letter
 LANGUAGE: English
 SUMMARY LANGUAGE: English

L5 ANSWER 8 OF 21 CAPLUS COPYRIGHT 2002 ACS
 ACCESSION NUMBER: 2000:227864 CAPLUS
 DOCUMENT NUMBER: 132:264105
 TITLE: YKL-40 as a marker and prognostic indicator for cancers
 INVENTOR(S): Price, Paul A.; Johansen, Julia S.
 PATENT ASSIGNEE(S): The Regents of the University of California, USA
 SOURCE: PCT Int. Appl., 111 pp. CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000019206	A1	20000406	WO 1999-US22615	19990929
W: AU, CA, JP, KR, NO				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
AU 9962757	A1	20000417	AU 1999-62757	19990929
EP 1112497	A1	20010704	EP 1999-950002	19990929
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				

PRIORITY APPLN. INFO.: US 1998-164862 A 19981001
 WO 1999-US22615 W 19990929

AB This invention provides methods for detecting cancers and for evaluating the prognosis of cancer patients. In particular, the methods of this invention utilize YKL-40 as a marker for the presence or absence of a cancer and for the prognosis (e.g. likelihood of recurrence) of a cancer. Elevated levels of YKL-40 are indicative of the presence of a cancer in undiagnosed subjects and indicate likely recurrence of the cancer in subjects diagnosed as having a cancer.

REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 9 OF 21 MEDLINE DUPLICATE 5
 ACCESSION NUMBER: 2001023517 MEDLINE
 DOCUMENT NUMBER: 20460900 PubMed ID: 11005786
 TITLE: Serum levels of YKL-40 and C reactive protein in patients with hip osteoarthritis and healthy subjects: a cross sectional study.
 AUTHOR: Conrozier T; Carlier M C; Mathieu P; Colson F; Debard A L; Richard S; Favret H; Bienvenu J; Vignon E
 CORPORATE SOURCE: Department of Rheumatology, Centre Hospitalier Lyon-Sud, Pierre-Benite, France.. th.conrozier@wanadoo.fr

SOURCE: ANNALS OF THE RHEUMATIC DISEASES, (2000 Oct) 59 (10)
828-31.
Journal code: 0372355. ISSN: 0003-4967.
PUB. COUNTRY: ENGLAND; United Kingdom
Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 200011
ENTRY DATE: Entered STN: 20010322
Last Updated on STN: 20010322
Entered Medline: 20001106

AB BACKGROUND: YKL-40 is a 40 kDa glycoprotein secreted by chondrocytes and synovial cells. It has been suggested that it is a surrogate marker of synovial inflammation and joint destruction in rheumatoid arthritis (RA) and osteoarthritis (OA) and related to C reactive protein (CRP) serum levels in RA. OBJECTIVE: To study serum levels of YKL-40 in patients with hip OA and its relation with CRP. METHODS: YKL-40 and CRP were assayed in serum samples from 45 patients (24 women, 21 men, mean age 65) with symptomatic OA of the hip and 33 healthy controls. YKL-40 was assayed by immunoassay and CRP by ultrasensitive immunonephelometry. OA severity was assessed by the measurement of joint space width with a computer analysis system of digitised hip radiographs. Statistical analysis was performed to determine correlations between serum markers and radiological joint space width. RESULTS: The mean (standard error) YKL-40 level was 90.3 (8.2) ng/ml in patients with hip OA and 66.9 (8.2) ng/ml in controls (p=0.03). The mean CRP level was 2.93 (3.03) mg/l in OA and 1.40 (1.61) mg/l in controls (p=0.006). The serum levels of YKL-40 and CRP increased with age and were significantly correlated (Spearman test: r(s)=0.42, p=0.005) in patients but not in controls. Neither YKL-40 nor CRP correlated with radiographic joint space width. CONCLUSIONS: Serum YKL-40 was significantly increased in patients with hip OA. The correlation between YKL-40 and CRP suggests that YKL-40 may be a marker of joint inflammation in OA. Longitudinal studies are required to assess the usefulness of YKL-40 in the monitoring of patients with hip OA.

L5 ANSWER 10 OF 21 MEDLINE DUPLICATE 6

ACCESSION NUMBER: 2000400313 MEDLINE
DOCUMENT NUMBER: 20334213 PubMed ID: 10873965
TITLE: Raised human cartilage glycoprotein-39 plasma levels in patients with rheumatoid arthritis and other inflammatory conditions.
AUTHOR: Vos K; Steenbakkers P; Miltenburg A M; Bos E; van Den Heuvel M W; van Hogeand R A; de Vries R R; Breedveld F C; Boots A M
CORPORATE SOURCE: Department of Rheumatology, LUMC, Leiden, The Netherlands.. kvos@rheumatology.azl.nl
SOURCE: ANNALS OF THE RHEUMATIC DISEASES, (2000 Jul) 59 (7) 544-8.
Journal code: 0372355. ISSN: 0003-4967.
PUB. COUNTRY: ENGLAND; United Kingdom
Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 200008
ENTRY DATE: Entered STN: 20000824
Last Updated on STN: 20000824
Entered Medline: 20000816

AB OBJECTIVE: To evaluate plasma human cartilage glycoprotein (HC gp-39) as a possible marker for the presence and/or activity of rheumatoid arthritis (RA) and other inflammatory conditions. BACKGROUND: HC gp-39 is a secretory product of chondrocytes, synovial cells, macrophages, and neutrophils. HC gp-39, also described as YKL-40, was found to be a marker of joint disease and tissue injury in RA and various other diseases. METHODS: Levels of HC gp-39 were determined by a sandwich enzyme linked immunosorbent assay (ELISA) in 47 patients with RA, 47 with osteoarthritis (OA), 24 with systemic lupus erythematosus (SLE), 24 with inflammatory bowel disease (IBD), and in 47 healthy controls. A disease activity score was assessed in the patients with RA, SLE, and IBD. RESULTS: The plasma level of HC gp-39 in the RA patient group was significantly higher than in the other patient groups and healthy controls. The level in patients with OA, SLE, and IBD was also significantly higher than the HC gp-39 level found in the healthy control group. HC gp-39 levels in patients with RA correlated positively with the ESR and IgM rheumatoid factor level but not with other variables of disease activity. In the patients with SLE and IBD no correlation was found with the disease activity score. CONCLUSION: The plasma level of HC gp-39 is increased in inflammatory conditions with and without joint disease (SLE, IBD, OA, and RA). Thus increased levels of HC gp-39 do not only reflect joint disease but also reflect inflammation or tissue degradation in various conditions. Notably, the highest level of HC gp-39 was found in patients with RA. Only in the RA patient group was a correlation between HC gp-39 plasma levels and some laboratory variables of disease activity found.

L5 ANSWER 11 OF 21 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: 2000:185155 BIOSIS
DOCUMENT NUMBER: PREV200000185155
TITLE: Serum chondrex values in knee osteoarthritis (OA). The effect of arthroscopy.
AUTHOR(S): Maciel, S. B.; Scheinberg, M. A. (1)
CORPORATE SOURCE: (1) 26 Dr Martinico Prado, Sao Paulo, SP, 01224-010 Brazil
SOURCE: Clinical Rheumatology, (2000) Vol. 19, No. 1, pp. 76-77. ISSN: 0770-3198.
DOCUMENT TYPE: Article
LANGUAGE: English
SUMMARY LANGUAGE: English

L5 ANSWER 12 OF 21 MEDLINE DUPLICATE 7

ACCESSION NUMBER: 2000488238 MEDLINE
DOCUMENT NUMBER: 20492231 PubMed ID: 11037631
TITLE: [Biochemical markers of bone turnover and YKL 40 in ankylosing spondylitis. Relation to disease activity]. Marcatori biochimici di turnover osseo e YKL 40 nella spondilite anchilosante. Rapporto con l'attivit  di malattia.
AUTHOR: D'Amore M; Germinario G; D'Amore S; Scagliusi P
CORPORATE SOURCE: Dipartimento di Medicina Interna e di Medicina Pubblica, Universita degli Studi, Bari.
SOURCE: MINERVA MEDICA, (2000 Mar-Apr) 91 (3-4) 59-68.
Journal code: 0400732. ISSN: 0026-4806.
PUB. COUNTRY: Italy

Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: Italian
FILE SEGMENT: Priority Journals
ENTRY MONTH: 200011
ENTRY DATE: Entered STN: 20010322
Last Updated on STN: 20010322
Entered Medline: 20001128

AB BACKGROUND: YKL-40 is a glycoprotein produced by chondrocytes and synovial cells. The plasmatic levels of this metabolite increase in some pathologies such as rheumatoid arthritis and osteoarthritis, so much so that it can be considered an effective marker of disease activity and in the therapeutic monitoring of these diseases. It has been interesting to dose a group of both male and female subjects affected by seronegative spondylarthritis, comparing this parameter with the disease activity indexes and with the bone turnover markers. METHODS: The study has been carried out on 48 subjects (26 males and 22 females) between 17 and 68 years affected by spondylarthritis, diagnosed in conformity with ARA standards. None of the patients carried out basic treatment or by glyocorticoids, and 22 patients took FANS when required. In these subjects the disease activity markers (VES, PCR, fibrinogen, mucoprotein) and some of the classic bone remodelling markers (blood calcium and phosphates, calciuria, phosphaturia, Ca++, Ntx, osteocalcine, bone isoenzyme of alkaline phosphatase, hydroxyproline, procollagen and YKL-40) were dosed. RESULTS: The comparison between different parameters pointed out that the highest values are obtained in subjects of most advanced age with the highest phlogosis indexes, without any correlation with sex. The quite interesting comparison shows a correlation between the bone remodelling indexes and YKL-40, being particularly remarkable when the disease is more aggressive or during relapse. CONCLUSIONS: It is then possible to confirm that, though preliminary, these data may suggest evaluations on wider case histories to research YKL-40 as a surgical monitoring marker in seronegative poliartthritis.

L5 ANSWER 13 OF 21 MEDLINE DUPLICATE 8

ACCESSION NUMBER: 1999390575 MEDLINE
DOCUMENT NUMBER: 99390575 PubMed ID: 10461474
TITLE: Serum YKL-40 concentrations in patients with rheumatoid arthritis: relation to disease activity.
AUTHOR: Johansen J S; Stoltenberg M; Hansen M; Florescu A; Horslev-Petersen K; Lorenzen I; Price P A
CORPORATE SOURCE: Department of Rheumatology, Hvidovre Hospital, Denmark.
SOURCE: RHEUMATOLOGY, (1999 Jul) 38 (7) 618-26.
JOURNAL code: 100883501. ISSN: 1462-0324.
PUB. COUNTRY: ENGLAND: United Kingdom
JOURNAL; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 199909
ENTRY DATE: Entered STN: 19990921
Last Updated on STN: 19990921
Entered Medline: 19990908

AB OBJECTIVE: YKL-40, also called human cartilage glycoprotein-39, is secreted by chondrocytes, synovial cells, macrophages and neutrophils. Studies have shown that YKL-40 is an autoantigen in rheumatoid arthritis (RA). We evaluated whether serum YKL-40 was related to disease activity in patients with RA. METHODS: Serum YKL-40 was determined by radioimmunoassay in 156 patients with RA during a 1 yr longitudinal study. RESULTS: Serum YKL-40 was increased in 54% of the patients with clinically active disease. Patients with clinically active disease initially who became inactive after 12 months had a significant decrease in serum YKL-40 (-30%, $P < 0.002$) and patients who changed from inactive to active disease had an increase in serum YKL-40. Patients who remained active had unchanged serum YKL-40 during the study. Serum YKL-40 decreased rapidly (-24% after 7 days, $P < 0.01$) during prednisolone therapy, and more slowly in patients treated with methotrexate only (-15% after 60 days, $P < 0.01$). Patients with early RA (disease duration < 3 yr, $n = 50$) and a persistently elevated serum YKL-40 were at risk of radiological disease progression as determined by Larsen score. CONCLUSION: Serum YKL-40 varies according to disease activity in RA, but provides in some respect information different from conventional markers. Our previous studies are consistent with a local release of YKL-40 in the arthritic joint followed by a secondary increase in serum YKL-40. YKL-40 may prove to be a new tool for the study of disease activity and pathophysiology of RA.

L5 ANSWER 14 OF 21 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.

ACCESSION NUMBER: 1999249149 EMBASE
TITLE: Nitric oxide alters chondrocyte function by disrupting cytoskeletal signaling complexes.
AUTHOR: Clancy R.
CORPORATE SOURCE: R. Clancy, Department of Rheumatology, Hospital for Joint Diseases, NYU School of Medicine, New York, NY, United States
SOURCE: Osteoarthritis and Cartilage, (1999) 7/4 (399-400).
Refs: 10
ISSN: 1063-4584 CODEN: OSCAEO
COUNTRY: United Kingdom
DOCUMENT TYPE: Journal; Conference Article
FILE SEGMENT: 029 Clinical Biochemistry
031 Arthritis and Rheumatism
LANGUAGE: English
SUMMARY LANGUAGE: English

AB Components of osteoarthritis include increases in pericellular fibronectin and in chondrocyte .beta.1 integrin expression. Events which follow ligation of fibronectin to its chondrocyte-receptor, the integrin .alpha.5.beta.1 include an assembly of a subplasmalemmal actin/rho A/focal adhesion kinase signaling complex. In addition, nitric oxide (NO), a potential mediator of cartilage pathophysiology disrupts the cytoskeletal signaling complex associated with integrin signaling. In these studies, we examined the relationship among integrin signaling, biosynthesis of S-35 sulfate containing proteoglycans and release of YKL-40 (a secretory glycoprotein) by comparing cell responses using cells plated on a fibronectin-coated or polyHEME coated surfaces. We report that the release of proteoglycan and glycoprotein require anchorage dependent signals by integrin costimulation. NO which disrupts the integrin signaling complex attenuates both cell responses. Taken together NO may serve as a nonspecific 'brake' to prevent anabolic and catabolic injury responses.

L5 ANSWER 15 OF 21 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: 1999:537094 BIOSIS

DOCUMENT NUMBER: PREV199900537094

TITLE: A novel arthritis model mice by secretory protein of articular chondrocytes, YKL-39.

AUTHOR(S): Sakata, Masahiro (1); Masuko-Hongo, Kayo; Tsuruha, Jun-ichiro; Nakamura, Hiroshi; Sekine, Taichi; Yoshino, Shin-ichi; Takigawa, Masaharu; Kato, Tomohiro; Nishioka, Kusuki

CORPORATE SOURCE: (1) Kawasaki Japan

SOURCE: Arthritis & Rheumatism, (Sept., 1999) Vol. 42, No. 9 SUPPL., pp. S257.

Meeting Info.: 63rd Annual Scientific Meeting of the American College of Rheumatology and the 34th Annual Scientific Meeting of the Association of Rheumatology Health Professionals Boston, Massachusetts, USA November 13-17, 1999

ISSN: 0004-3591.

DOCUMENT TYPE: Conference

LANGUAGE: English

L5 ANSWER 16 OF 21 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: 1999:537080 BIOSIS

DOCUMENT NUMBER: PREV199900537080

TITLE: Regulation of YKL-40 production by human articular chondrocytes.

AUTHOR(S): Johansen, Julia S. (1); Olee, Tsaiwei (1); Price, Paul A. (1); Ochs, Robert L. (1); Hashimoto, Sanshiro (1); Lotz, Martin (1)

CORPORATE SOURCE: (1) La Jolla, CA USA

SOURCE: Arthritis & Rheumatism, (Sept., 1999) Vol. 42, No. 9 SUPPL., pp. S255.

Meeting Info.: 63rd Annual Scientific Meeting of the American College of Rheumatology and the 34th Annual Scientific Meeting of the Association of Rheumatology Health Professionals Boston, Massachusetts, USA November 13-17, 1999

ISSN: 0004-3591.

DOCUMENT TYPE: Conference

LANGUAGE: English

L5 ANSWER 17 OF 21 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: 1999:535492 BIOSIS

DOCUMENT NUMBER: PREV199900535492

TITLE: Serum YKL-40, a chondrocyte derived protein is reduced by infliximab (anti-TNF) therapy in patients with rheumatoid arthritis.

AUTHOR(S): Charles, P. J. (1); Maini, R. N. (1)

CORPORATE SOURCE: (1) London UK

SOURCE: Arthritis & Rheumatism, (Sept., 1999) Vol. 42, No. 9 SUPPL., pp. S236.

Meeting Info.: 63rd Annual Scientific Meeting of the American College of Rheumatology and the 34th Annual Scientific Meeting of the Association of Rheumatology Health Professionals Boston, Massachusetts, USA November 13-17, 1999

ISSN: 0004-3591.

DOCUMENT TYPE: Conference

LANGUAGE: English

L5 ANSWER 18 OF 21 MEDLINE DUPLICATE 9

ACCESSION NUMBER: 1999308547 MEDLINE

DOCUMENT NUMBER: 99308547 PubMed ID: 10380840

TITLE: The distribution of YKL-40 in osteoarthritic and normal human articular cartilage.

AUTHOR: Volck B; Ostergaard K; Johansen J S; Garbarsch C; Price P A

CORPORATE SOURCE: Department of Rheumatology, Hvidovre Hospital, Denmark.

SOURCE: SCANDINAVIAN JOURNAL OF RHEUMATOLOGY, (1999) 28 (3) 171-9.

Journal code: 0321213. ISSN: 0300-9742.

PUB. COUNTRY: Norway

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 199906

ENTRY DATE: Entered STN: 19990714

Last Updated on STN: 19990714

Entered Medline: 19990629

AB YKL-40, also called human cartilage glycoprotein-39, is a major secretory protein of human chondrocytes in cell culture. YKL-40 mRNA is expressed by cartilage from patients with rheumatoid arthritis, but is not detectable in normal human cartilage. The aim was to investigate the distribution of YKL-40 in osteoarthritic (n=9) and macroscopically normal (n=5) human articular cartilage, collected from 12 pre-selected areas of the femoral head, to discover a potential role for YKL-40 in cartilage remodelling in osteoarthritis. Immunohistochemical analysis showed that YKL-40 staining was found in chondrocytes of osteoarthritic cartilage mainly in the superficial and middle zone of the cartilage rather than the deep zone. There was a tendency for high number of YKL-40 positive chondrocytes in areas of the femoral head with a considerable biomechanical load. The number of chondrocytes with a positive staining for YKL-40 was in general low in normal cartilage. The present findings, together with previous observations, suggests that YKL-40 may be of importance in cartilage remodelling/degradation of osteoarthritic joints.

L5 ANSWER 19 OF 21 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1998:170374 CAPLUS

DOCUMENT NUMBER: 128:280523

TITLE: Chondrex: new marker of joint disease

AUTHOR(S): Harvey, Sheryl; Weisman, Michael; O'Dell, James; Scott, Tonya; Krusemeier, Mindy; Visor, Jill; Swindlehurst, Cathy

CORPORATE SOURCE: Novadex, Inc, San Diego, CA, 92121, USA

SOURCE: Clinical Chemistry (Washington, D. C.) (1998), 44(3), 509-516

CODEN: CLCHAU; ISSN: 0009-9147

PUBLISHER: American Association for Clinical Chemistry

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Chondrex, a major secretory protein of human chondrocytes and synovial fibroblasts, is increased in serum of patients with joint and cartilage disease. We have developed a sandwich-type ELISA for

quantifying chondrex in serum. The interassay CVs were 2.8-3.7% and the av. within-run and total CVs were 3.6% and 5.4%, resp. The limit of detectability by linear diln. was 20 .mu.g/L, recovery upon diln. was 102% +/- 5%, and anal. recovery (of added analyte) was 98% +/- 11%. The ref. interval (central 90% interval) for chondrex in healthy adults was 25-95 .mu.g/L. Chondrex values for patients with active rheumatoid arthritis or osteoarthritis were significantly greater than in healthy adults, inactive rheumatoid arthritis patients, and diabetes patients ($P < 0.05$). In patients treated with disease-modifying antirheumatic drug therapy, decreasing chondrex values reflected the clin. improvement obsd. in responders, whereas the values were maintained or increased in nonresponders. In conclusion, chondrex may be a useful marker in the clin. investigation of arthritis.

L5 ANSWER 20 OF 21 MEDLINE DUPLICATE 10
 ACCESSION NUMBER: 1998349472 MEDLINE
 DOCUMENT NUMBER: 98349472 PubMed ID: 9686683
 TITLE: YKL-40, a mammalian member of the chitinase family, is a matrix protein of specific granules in human neutrophils.
 AUTHOR: Volck B; Price P A; Johansen J S; Sorensen O; Benfield T L; Nielsen H J; Calafat J; Borregaard N
 CORPORATE SOURCE: Department of Rheumatology, Hvidovre Hospital, University of Copenhagen, Denmark.
 SOURCE: PROCEEDINGS OF THE ASSOCIATION OF AMERICAN PHYSICIANS, (1998 Jul-Aug) 110 (4) 351-60.
 PUB. COUNTRY: United States
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 199904
 ENTRY DATE: Entered STN: 19990426
 Last Updated on STN: 19990426
 Entered Medline: 19990413

AB YKL-40, also called human cartilage glycoprotein-39 (HC gp-39), is a member of family 18 glycosyl hydrolases. YKL-40 is secreted by chondrocytes, synovial cells, and macrophages, and recently it has been reported that YKL-40 has a role as an autoantigen in rheumatoid arthritis (RA). The function of YKL-40 is unknown, but the pattern of its expression in normal and disease states suggests that it could function in remodeling or degradation of the extracellular matrix. High levels of YKL-40 are found in synovial fluid from patients with active RA. Neutrophils are abundant in synovial fluid of patients with RA, and the cells are assumed to play a role in joint destruction in that disorder. Therefore, we examined whether neutrophils are a source of YKL-40. YKL-40 was found to colocalize and comobilize with lactoferrin (the most abundant protein of specific granules) but not with gelatinase in subcellular fractionation studies on stimulated and unstimulated neutrophils. Double-labeling immunoelectron microscopy confirmed the colocalization of YKL-40 and lactoferrin in specific granules of neutrophils. Immunohistochemistry on bone marrow cells showed that neutrophil precursors begin to synthesize YKL-40 at the myelocyte-metamyelocyte stage, the stage of maturation at which other specific granule proteins are formed. Assuming that YKL-40 has a role as an autoantigen in RA by inducing T cell-mediated autoimmune response, YKL-40 released from neutrophils in the inflamed joint could be essential for this response. In RA and other inflammatory diseases, YKL-40 released from specific granules of neutrophils may be involved in tissue remodeling or degradation.

L5 ANSWER 21 OF 21 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.
 ACCESSION NUMBER: 93334909 EMBASE
 DOCUMENT NUMBER: 1993334909
 TITLE: A new biochemical marker for joint injury. Analysis of YKL 40 in serum and synovial fluid.
 AUTHOR: Johansen J.S.; Jensen H.S.; Price P.A.
 CORPORATE SOURCE: Dept of Med, Div of Rheumatology 232, University of Copenhagen, Hvidovre Hospital, Kettegaard Alle 30, Hvidovre DK-2650, Denmark
 SOURCE: British Journal of Rheumatology, (1993) 32/11 (949-955).
 COUNTRY: United Kingdom
 DOCUMENT TYPE: Journal; Article
 FILE SEGMENT: 031 Arthritis and Rheumatism
 LANGUAGE: English
 SUMMARY LANGUAGE: English

AB We report the development of the first radioimmunoassay for YKL-40, a M(r) = 40 kDa protein which is secreted at high levels by human synovial cells and articular cartilage chondrocytes, and by the human osteosarcoma cell line MG63. This assay uses YKL-40 purified from the conditioned medium of MG63 cells as standard and tracer, and as antigen for immunizing rabbits. With this assay we have discovered high levels of YKL-40 antigen in serum and SF. The molecular weight of serum and SF YKL-40 is identical to purified YKL-40. To evaluate the possible utility of YKL-40 in the assessment of joint disease, we measured YKL-40 in serum and SF of 49 patients with various forms of inflammatory and degenerative joint disease and in the serum of 50 normal adults. The YKL-40 level in serum was significantly higher ($P < 0.001$) in the patients compared to the normal adults, but there was no difference in serum YKL-40 between the patients with inflammatory joint diseases and OA. The SF levels of YKL-40 were 15-fold higher than serum levels and there was a significant correlation ($r = 0.55$, $P < 0.001$) between YKL-40 concentration in SF and serum. Although the tissue distribution of YKL-40 secretion is presently unknown, these observations suggest that a major portion of serum YKL-40 in fact arises from the joint. Serum and SF YKL 40 levels are correlated significantly ($P < 0.05$ - $P < 0.001$) with other indices of joint disease: serum CRP, SF IL-6, and the elastolytic activity of monocytes/macrophages in SF. Serum YKL-40 also correlated with serum PIIINP and elastolytic activity of blood monocytes/macrophages. These studies indicate that serum and SF YKL-40 levels reflect joint disease and a YKL-40 determination may therefore be useful in the evaluation of connective tissue injury and repair in patients with inflammatory or degenerative rheumatic diseases. Future studies will be needed in order to assess the physiologic significance of elevated YKL-40 levels in patients with rheumatoid diseases.

L6 19 YKL (1N) 39

=> dup rem 16

PROCESSING COMPLETED FOR L6

L7 10 DUP REM L6 (9 DUPLICATES REMOVED)

=> dis 17 1-10 ibib abs

L7 ANSWER 1 OF 10 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: 2002:261273 BIOSIS

DOCUMENT NUMBER: PREV200200261273

TITLE: Analysis of chondrex (YKL-40, HC gp-39) in the cerebrospinal fluid of patients with spine disease.

AUTHOR(S): Tsuji, Taichi (1); Matsuyama, Yukihiro; Natsume, Naoki; Hasegawa, Yukiharu; Kondo, Seiji; Kawakami, Hiroshi; Yoshihara, Hisatake; Iwata, Hisashi

CORPORATE SOURCE: (1) Department of Orthopaedic Surgery, Nagoya University School of Medicine, 65 Tsuruma-cho, Showa-ku, Nagoya, 466-8550; tsuji-t@med.nagoya-u.ac.jp Japan

SOURCE: Spine, (April 1, 2002) Vol. 27, No. 7, pp. 732-735.

http://www.spinejournal.com/. print.

ISSN: 0362-2436.

DOCUMENT TYPE: Article

LANGUAGE: English

AB Study Design: The expression of chondrex (YKL-40, HC gp-39) was measured in the cerebrospinal fluid of from patients with spine diseases. Objectives: To quantify the levels of chondrex in human cerebrospinal fluid, and to clarify the nature of its expression. Summary of Background Data: Chondrex is a newly discovered 40-kDa glycoprotein identified originally in the whey secretions of nonlactating cows. It is secreted by a human osteosarcoma cell line, human articular cartilage chondrocytes, and human fibroblasts. However, the function of chondrex in chondrogenesis is unknown, and the expression of chondrex in human cerebrospinal fluid has never been reported. Methods: The concentration of chondrex in human cerebrospinal fluid was measured by sandwich immunoassay with antihuman chondrex antibodies. Cerebrospinal fluid samples were collected from two groups of patients. Group 1, the control group, consisted of 34 trauma patients. Group 2 consisted of 130 patients with spine diseases: 29 with cervical spondylotic myelopathy, 30 with lumbar disc herniation, 35 with lumbar canal stenosis, and 36 with scoliosis. All values are expressed as the mean±standard deviation. Results: The concentration of chondrex in Group 1 (control group) was 113.8±48.3 ng/mL. The concentrations of chondrex in Group 2 were 245.3±107.2 ng/mL in cervical myelopathy, 143.2±53.6 ng/mL in lumbar disc herniation, 241.5±77.2 ng/mL in lumbar canal stenosis, and 71.4±33.9 ng/mL in scoliosis. The concentrations of chondrex in cervical myelopathy, lumbar canal stenosis, and lumbar disc herniation were significantly higher than in the control group (P<0.05). Conclusions: In this study, the chondrex concentration was high in spine diseases causing spinal stenosis. The authors believe that chondrex is expressed in cerebrospinal fluid as a result of damage or stress to the neural structure, and that it could be a new marker for spine diseases.

L7 ANSWER 2 OF 10

MEDLINE

DUPLICATE 1

ACCESSION NUMBER: 2001252197 MEDLINE

DOCUMENT NUMBER: 21248333 PubMed ID: 11350852

TITLE: Cross sectional evaluation of biochemical markers of bone, cartilage, and synovial tissue metabolism in patients with knee osteoarthritis: relations with disease activity and joint damage.

COMMENT: Comment in: Ann Rheum Dis. 2001 Jun;60(6):545-8

AUTHOR: Garnero P; Piperno M; Gineyts E; Christgau S; Delmas P D; Vignon E

CORPORATE SOURCE: Inserm Research Unit 403, Lyon, France..

patrick.garnero@synarc.com

SOURCE: ANNALS OF THE RHEUMATIC DISEASES, (2001 Jun) 60 (6) 619-26.

Journal code: 0372355. ISSN: 0003-4967.

PUB. COUNTRY: England: United Kingdom

(EVALUATION STUDIES)

Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 200106

ENTRY DATE: Entered STN: 20010611

Last Updated on STN: 20010611

Entered Medline: 20010607

AB OBJECTIVE: To analyse the relations between the urinary levels of type II collagen C-telopeptide (CTX-II) and glucosyl-galactosyl pyridinoline (Glc-Gal-PYD)-two newly developed biochemical markers of type II collagen and synovial tissue destruction respectively-disease activity and the severity of joint destruction in patients with knee osteoarthritis (OA). The clinical performance of these two new markers was compared with that of a panel of other established biochemical markers of connective tissue metabolism. METHODS: The following biochemical markers were measured in a group of 67 patients with knee OA (mean age 64 years, median disease duration eight years) and in 67 healthy controls: for bone, serum osteocalcin, serum and urinary C-telopeptide of type I collagen (CTX-I); for cartilage, urinary CTX-II, serum cartilage oligomeric matrix protein (COMP), and serum human cartilage glycoprotein 39 (YKL-40); for synovium, urinary Glc-Gal-PYD, serum type III collagen N-propeptide (PIIINP), serum hyaluronic acid (HA); and for inflammation, serum C reactive protein. Biochemical markers were correlated with pain and physical function (WOMAC index) and with quantitative radiographic evaluation of the joint space using the posteroanterior view of the knees flexed at 30 degrees. RESULTS: All bone turnover markers were decreased in patients with knee OA compared with controls (-36%, -38%, and -52%, p<0.0001 for serum osteocalcin, serum CTX-I and urinary CTX-I, respectively). Serum COMP (+16%, p=0.0004), urinary CTX-II (+25%, p=0.0009), urinary Glc-Gal-PYD (+18%, p=0.028), serum PIIINP (+33%, p<0.0001), and serum HA (+233%, p<0.0001) were increased. By univariate analyses, increased urinary Glc-Gal-PYD (r=0.41, p=0.002) and decreased serum osteocalcin (r=-0.30, p=0.025) were associated with a higher total WOMAC index. Increased urinary CTX-II (r=-0.40, p=0.0002) and Glc-Gal-PYD (r=-0.30, p=0.0046) and serum PIIINP (r=-0.29, p=0.0034) were the only markers which correlated with joint surface area. By multivariate analyses, urinary Glc-Gal-PYD and CTX-II were the most important predictors of the WOMAC index and joint damage, respectively. CONCLUSION: Knee OA appears to be characterised by a systemic decrease of bone turnover and increased cartilage and synovial tissue turnover. CTX-II, Glc-Gal-PYD, and PIIINP may be useful markers of disease severity in patients with knee OA.

L7 ANSWER 3 OF 10 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 2001:315060 CAPLUS

DOCUMENT NUMBER: 134:363566
 TITLE: Development of an enzyme-linked immunoassay for the quantification of YKL-40 (cartilage gp-39) in guinea pig serum using hen egg yolk antibodies
 AUTHOR(S): De Ceuninck, P.; Pastoureau, P.; Agnellet, S.; Bonnet, J.; Vanhoutte, P. M.
 CORPORATE SOURCE: Division of Rheumatology, Institut de Recherches Servier, Suresnes, 92150, Fr.
 SOURCE: Journal of Immunological Methods (2001), 252(1-2), 153-161
 CODEN: JIMMBG; ISSN: 0022-1759
 PUBLISHER: Elsevier Science B.V.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB An indirect competition immunoassay for the quantification of YKL-40 (cartilage gp-39, Chondrex) in guinea pig serum has been developed using egg yolk antibodies (IgY). The immune response of hens to YKL-40 was verified by immunoblot analyses. Highly specific antibodies were obtained 30 days after the first injection. The ELISA was developed in 96-well microtiter plates with quadruplicate detns. for each point. The assay was based on the ability of YKL-40 present in serum to displace the binding of antibodies to the coated antigen. An inhibition mixt. contg. std. YKL-40 or guinea pig serum, dild. 1/5, and primary antibodies, dild. 1/5000, was allowed to equilibrate for 2 h at room temp. and dispensed for 16 h at 4.degree. in wells coated with 1 .mu.g/mL of YKL-40. Detection was achieved by the addn. of rabbit anti-chicken antibodies conjugated to peroxidase followed by tetramethylbenzidine. Specificity was assessed by parallelism between a diln. curve of serum and std. YKL-40. The sensitivity of detection was 10 ng/mL. Intra- and interassay coeffs. of variation were both 8.7%. The anal. recovery was 101.5+-5.4% (mean+-std. deviation (SD), n=9). The YKL-40 concn. in serum from 12 adult guinea pigs was 330+-216 ng/mL (mean+-SD) with a lower value of 164 ng/mL and an upper value of 982 ng/mL. In contrast to the rat, a diln. curve of rabbit serum gave parallelism with the guinea pig std., suggesting recognition of a similar epitope. Possible applications of the assay in the guinea pig include disease models where YKL-40 is overexpressed and could be used as a marker, i.e. osteoarthritis, rheumatoid arthritis, cancer, liver fibrosis, atherosclerosis and more generally, pathologies with increased tissue remodeling.
 REFERENCE COUNT: 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 4 OF 10 MEDLINE DUPLICATE 2
 ACCESSION NUMBER: 2001124047 MEDLINE
 DOCUMENT NUMBER: 20566402 PubMed ID: 11114282
 TITLE: Recognition of YKL-39, a human cartilage related protein, as a target antigen in patients with rheumatoid arthritis.
 AUTHOR: Sekine T; Masuko-Hongo K; Matsui T; Asahara H; Takigawa M; Nishioka K; Kato T
 CORPORATE SOURCE: Rheumatology, Immunology and Genetics Programme, Institute of Medical Science, St Marianna University, School of Medicine 2-16-1, Sugao, Miyamae-ku, Kawasaki, Kanagawa, 216-8512, Japan.
 SOURCE: ANNALS OF THE RHEUMATIC DISEASES, (2001 Jan) 60 (1) 49-54.
 PUB. COUNTRY: ENGLAND: United Kingdom
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 200102
 ENTRY DATE: Entered STN: 20010322
 Last Updated on STN: 20010322
 Entered Medline: 20010222
 AB OBJECTIVE: To investigate whether autoimmunity to YKL-39, a recently cloned cartilage protein, occurs in patients with rheumatoid arthritis (RA). METHODS: Autoantibody to YKL-39 was assayed by enzyme linked immunosorbent assay (ELISA) and western blotting in serum samples from patients with RA, systemic lupus erythematosus (SLE), and healthy donors, using recombinant YKL-39 protein. This reactivity was compared with that against a YKL-39 homologue, YKL-40 (human cartilage gp-39/chondrex), which has been reported to be an autoantigen in RA. RESULTS: Autoantibody to YKL-39 was detected in seven of 87 patients with RA (8%), but not in serum samples from patients with SLE or healthy donors. YKL-40 reactivity was found in only one of 87 RA serum samples (1%), with no cross reactivity to YKL-39. CONCLUSION: The existence of anti-YKL-39 antibody in a subset of patients with RA is reported here for the first time. Further, it was shown that the immune response to YKL-39 was independent of that to YKL-40. Clarification of the antibody and T cell responses to autoantigens derived from chondrocyte, cartilage, or other joint components may lead to a better understanding of the pathophysiology of joint destruction in patients with RA.

L7 ANSWER 5 OF 10 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
 ACCESSION NUMBER: 1999:537094 BIOSIS
 DOCUMENT NUMBER: PREV199900537094
 TITLE: A novel arthritis model mice by secretory protein of articular chondrocytes, YKL-39.
 AUTHOR(S): Sakata, Masahiro (1); Masuko-Hongo, Kayo; Tsuruha, Jun-ichiro; Nakamura, Hiroshi; Sekine, Taichi; Yoshino, Shin-ichi; Takigawa, Masaharu; Kato, Tomohiro; Nishioka, Kusuiki
 CORPORATE SOURCE: (1) Kawasaki Japan
 SOURCE: Arthritis & Rheumatism, (Sept., 1999) Vol. 42, No. 9 SUPPL., pp. S257.
 Meeting Info.: 63rd Annual Scientific Meeting of the American College of Rheumatology and the 34th Annual Scientific Meeting of the Association of Rheumatology Health Professionals Boston, Massachusetts, USA November 13-17, 1999
 ISSN: 0004-3591.
 DOCUMENT TYPE: Conference
 LANGUAGE: English

L7 ANSWER 6 OF 10 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
 ACCESSION NUMBER: 1999:537097 BIOSIS
 DOCUMENT NUMBER: PREV199900537097
 TITLE: Argument of articular immune responses to gp-39 and YKL-39 in patients with osteoarthritis.
 AUTHOR(S): Tsuruha, Jun-ichiro (1); Masuko-Hongo, Kayo; Sakata,

Masahiro; Nakamura, Hiroshi; Sekine, Taichi; Yoshino, Shin-ichi; Takigawa, Masaharu; Kato, Tomohiro; Nishioka, Kusuki
 CORPORATE SOURCE: (1) Kawasaki Japan
 SOURCE: Arthritis & Rheumatism, (Sept., 1999) Vol. 42, No. 9 SUPPL., pp. S257.
 Meeting Info.: 63rd Annual Scientific Meeting of the American College of Rheumatology and the 34th Annual Scientific Meeting of the Association of Rheumatology Health Professionals Boston, Massachusetts, USA November 13-17, 1999
 ISSN: 0004-3591.
 DOCUMENT TYPE: Conference
 LANGUAGE: English

L7 ANSWER 7 OF 10 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.
 ACCESSION NUMBER: 1999:432310 BIOSIS
 DOCUMENT NUMBER: PREV199900432310
 TITLE: Possible quantitative trait loci for serum levels of human cartilage glycoprotein-39 (YKL-40) and osteocalcin (OC) in pedigree baboons map to human chromosomes 6 and 12.
 AUTHOR(S): Mahaney, M. C. (1); Czerwinski, S. A. (1); Rogers, J. (1)
 CORPORATE SOURCE: (1) Genetics, Southwest Foundation for Biomedical Research, San Antonio, TX USA
 SOURCE: Journal of Bone and Mineral Research, (Sept., 1999) Vol. 14, No. SUPPL. 1, pp. S142.
 Meeting Info.: Twenty-First Annual Meeting of the American Society for Bone and Mineral Research St. Louis, Missouri, USA September 30-October 4, 1999 American Society for Bone and Mineral Research
 . ISSN: 0884-0431.
 DOCUMENT TYPE: Conference
 LANGUAGE: English

L7 ANSWER 8 OF 10 CAPLUS COPYRIGHT 2002 ACS
 ACCESSION NUMBER: 1997:718004 CAPLUS
 DOCUMENT NUMBER: 128:16403
 TITLE: Human cartilage autoantigen glycoprotein gp-39 and proteins structurally related thereto for use in immunotherapy of autoimmune diseases
 INVENTOR(S): Boots, Anna Maria Helena; Verheijden, Gijsbertus Franciscus Maria; Bos, Ebo Sybren
 PATENT ASSIGNEE(S): Akzo Nobel N.V., Neth.; Boots, Anna Maria Helena; Verheijden, Gijsbertus Franciscus Maria; Bos, Ebo Sybren
 SOURCE: PCT Int. Appl., 38 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 3
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9740149	A1	19971030	WO 1997-EP1903	19970415
W: AM, AU, BB, BG, BR, CA, CN, CZ, EE, GE, HU, IS, JP, KG, KP, KR, LK, LR, LT, LV, MD, MG, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TR, TT, UA, US, UZ, VN, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
US 5843449	A	19981201	US 1996-634493	19960418
AU 9723869	A1	19971112	AU 1997-23869	19970415
AU 724547	B2	20000928		
EP 904369	A1	19990331	EP 1997-919370	19970415
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
BR 9708714	A	19990803	BR 1997-8714	19970415
JP 2000509265	T2	20000725	JP 1997-537477	19970415
NO 9804835	A	19981216	NO 1998-4835	19981016
PRIORITY APPLN. INFO.: US 1996-634493 A 19960418				
US 1996-619645 A2 19960325				
WO 1997-EP1903 W 19970415				

AB The present invention relates to the use of autoantigen HC gp-39 (human cartilage glycoprotein-39), and proteins comprising an amino acid sequence which exhibits .gtoreq.50% homol. with the amino acid sequence of HC gp-39, and more particular with the amino acid sequence YKLVCYITWSQYREGDGSFPDADRFLCTHIYSPANISND in antigen-specific treatment of articular cartilage destruction in autoimmune diseases in mammals to induce systemic tolerance of the immune system. The autoantigen HC gp-39, and the arthritogenic proteins comprising an amino acid sequence which exhibits .gtoreq.50% homol. with the amino acid sequence YKLVCYITWSQYREGDGSFPDADRFLCTHIYSPANISND are also suitable to induce arthritis in animals, preferably mice. The arthritogenicity and cloning of bovine 39-kDa whey protein is also described. The invention furthermore relates to pharmaceutical compns. comprising said autoantigen and/or said arthritogenic proteins, a diagnostic method for the detection of autoreactive T cells in a test sample and test kits to be used in said method.

L7 ANSWER 9 OF 10 CAPLUS COPYRIGHT 2002 ACS
 ACCESSION NUMBER: 1997:383815 CAPLUS
 DOCUMENT NUMBER: 127:62133
 TITLE: Various roles of chitinases
 AUTHOR(S): Watanabe, Takeshi
 CORPORATE SOURCE: Nogakubu, Niigata Daigaku, Niigata, 950-21, Japan
 SOURCE: Kagaku to Seibutsu (1997), 35(6), 408-414
 CODEN: KASEAA; ISSN: 0453-073X
 PUBLISHER: Gakkai Shuppan Senta
 DOCUMENT TYPE: Journal; General Review
 LANGUAGE: Japanese

AB A review, with 15 refs., on occurrence, physiol. function, classification, and conformation of chitinases. Reaction mechanism of chitinases of family 18 and proteins structurally analogous to family 18 chitinases, e.g. cartilage glycoprotein gp-39 (YKL-40), oviductin, concanavalin B, narbonin. etc., are also discussed.

L7 ANSWER 10 OF 10 MEDLINE DUPLICATE 3
 ACCESSION NUMBER: 96325055 MEDLINE
 DOCUMENT NUMBER: 96325055 PubMed ID: 8702629
 TITLE: Isolation and sequence of a novel human chondrocyte protein related to mammalian members of the chitinase protein

family.
 AUTHOR: Hu B; Trinh K; Figueira W P; Price P A
 CORPORATE SOURCE: Department of Biology, University of California, San Diego,
 La Jolla, California 92093, USA.
 CONTRACT NUMBER: AG07996 (NIA)
 SOURCE: JOURNAL OF BIOLOGICAL CHEMISTRY, (1996 Aug 9) 271 (32)
 19415-20.
 Journal code: 2985121R. ISSN: 0021-9258.
 PUB. COUNTRY: United States
 Journal; Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 OTHER SOURCE: GENBANK-U49835
 ENTRY MONTH: 199609
 ENTRY DATE: Entered STN: 19960924
 Last Updated on STN: 20010919
 Entered Medline: 19960916

AB We describe the isolation of a novel protein from the conditioned medium of human articular cartilage chondrocytes in primary culture. This 39-kDa protein has the N-terminal sequence YKL, which we have termed YKL-39. The 1434-nucleotide sequence of the YKL-39 cDNA predicts a 385-residue initial translation product and a 364-residue mature YKL-39. The amino acid sequence of YKL-39 is most closely related to YKL-40, followed by macrophage chitotriosidase, oviductal glycoprotein, and macrophage YM-1. All five proteins share significant sequence identity with bacterial chitinases and have the probable structure of an (alphabeta)8 barrel. YKL-39 lacks the active site glutamate, which is essential for the activity of chitinases, and as expected has no chitinase activity. The highest level of YKL-39 mRNA expression is seen in chondrocytes, followed by synoviocytes, lung, and heart. YKL-39 accounts for 4% of the protein in chondrocyte-conditioned medium, prostromelysin accounts for 17%, and YKL-40 accounts for 33%. In contrast to YKL-40, YKL-39 is not a glycoprotein and does not bind to heparin.

=> dis his

(FILE 'HOME' ENTERED AT 09:46:22 ON 25 JUN 2002)

FILE 'MEDLINE, CAPLUS, EMBASE, BIOSIS' ENTERED AT 09:46:36 ON 25 JUN 2002

L1 0 S FILASAEIT OR HSFILASAEITVG
 L2 211 S YKL
 L3 69 S L2 AND CHONDROCYTE?
 L4 42 S L3 AND (AUTOIMMUN? OR RA OR ARTHRITIS)
 L5 21 DUP REM L4 (21 DUPLICATES REMOVED)
 L6 19 S YKL (1N) 39
 L7 10 DUP REM L6 (9 DUPLICATES REMOVED)

=> end

ALL L# QUERIES AND ANSWER SETS ARE DELETED AT LOGOFF

LOGOFF? (Y)/N/HOLD:y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	60.61	60.82

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-3.10	-3.10

STN INTERNATIONAL LOGOFF AT 09:54:41 ON 25 JUN 2002